

Possible Single Nucleotide Polymorphism (SNP) in the Nucleic Sequence of A-kinase-anchoring Protein 9

Viroj Wiwanitkit

Wiwanitkit House, Bangkhae Bangkok Thailand 10160

Corresponding author: Viroj Wiwanitkit, M.D. Wiwanitkit House,
38/167 Soi Yim Prayoon, Bangkhae Bangkok Thailand 1016
Email: wviroj@yahoo.com

Received June 25, 2008; Accepted July 16, 2008; Published July 17, 2008

Citation: Viroj W (2008) Possible Single Nucleotide Polymorphism (SNP) in the Nucleic Sequence of A-kinase-anchoring Protein 9. *J Proteomics Bioinform* 1: 227-229.

Copyright: © 2008 Viroj W. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

In cardiac physiology, ion channel macromolecular complexes can be formed by the strong correlation of A-kinase anchoring proteins, therefore, AKAP becomes an important antiarrhythmic target. Here, the author performed a bioinformatics analysis to study the possible single nucleotide polymorphism (SNP) in the amino acid sequence of AKAP9. In this work, the author could not identify any SNP within AKAP9. This confirms that there is no report on SNP on this molecule.

Key words: A-kinase anchoring protein; arrhythmia; single nucleotide polymorphism

Introduction

In normal cardiac physiology, sympathetic nervous system (SNS) regulation of cardiac action potential duration (APD), mediated by beta-adrenergic receptor (betaAR) activation, usually requires assembly of A-kinase anchoring protein (AKAP)9 (Yotiao) with the I(Ks) potassium channel alpha subunit (KCNQ1) (Chen et al., 2001). Ion channel macromolecular complexes can be formed by the strong correlation of AKAPs (Chen et al., 2006, Marx and Kurokawa, 2006). Therefore, AKAP becomes an important antiarrhythmic target (Marx and Kurokawa, 2006). Analysis on the AKAP in depth helps better understand the pathogenesis of this protein-related arrhythmic disorder. AKAP disorder can be seen in the recent reports (Lin et al., 1998, Tao et al., 2006), however, the prevalence of mutations in the AKAP is not well known. In addition, single nucleotide polymorphism (SNP) within AKAP has never been mentioned.

Importance of SNP study for AKAP and other AKAP targets for new therapeutic interventions should be mentioned. Burns-Hamuro et al. said that there had been considerable progress in understanding the structural features of this AKAP and its interaction with protein kinase A (Burns-Hamuro et al., 2004). Burns-Hamuro et al. reported that comprehensive analysis of the PKA binding motif could lead to the development of novel peptides derived from D-AKAP2 and could be useful tools in probing the function of

this AKAP in cellular and animal models (Burns-Hamuro et al., 2004). However, there is limited reported on AKAP9. There was a paper by Rudd et al. describing the possible correlation of SNP for AKAP9 and lung cancer (Rudd et al., 2006). However, there was no proof from this work (Rudd et al., 2006). Here, the author performed a bioinformatics analysis to study the SNP in the amino acid sequence of AKAP9.

Materials and Methods

At first, the database Expert Protein Analysis System (ExPASy) (Gasteiger et al., 2003) was used for searching for the amino acid sequence of AKAP9. Then the derived sequences were used for further study for SNP. GEN-SNiP was further used for identification on SNP within the derived sequence. Briefly, GEN-SNiP helps find polymorphisms present in DNA sequences with respect to a standard reference sequence. This tool is selected because it is only one simple online assessment tool and is confirmed for its validity. The output lists substitutions, insertions and deletions. Since there is no other similar tool to GEN-SNiP, further comparative study to other tool cannot be performed.

Result

In this work, AKAP9 (NG_007968) was used for further study. There is no identified SNP for this sequence.

Discussion

Importance of SNP search for target drug discovery/new therapeutic interventions should be mentioned in the present post-genomics era. The analysis of SNPs permits determining relationships between genotypic and phenotypic information as well as the identification of SNPs related to a disease, which can be useful for further drug development (Shah and Kusiak, 2004). The good examples are on antineoplastic (Mack et al., 2008) and cardiovascular drugs (Borgiani et al., 2007). There are numerous experimental and clinical data supporting the existence or variant AKAP proteins (Chen et al., 2001). However, the exact functions of variant AKAP proteins, either physiological or pathological, are still unclear, although roles for some AKAP variants in cardiac arrhythmia progression might be consistent with the accumulated data (Chen et al., 2001; Saucerman et al., 2004). On one hand, AKAP polymorphism might connect with some rare syndromes of arrhythmic disorder and on other side it is the post-receptor modifications that very well explain the molecular pathogenesis of AKAP and arrhythmia. Identification of the SNP within AKAP can be useful for further researches to understand the pathogenesis of AKAP disorder. In this work, the author could not identify any SNP within AKAP9. This confirms that there is no report on SNP on this molecule although there is for other AKAPs ((Burns-Hamuro et al., 2004; Frank et al., 2004).

Conclusions & Future Perspectives

In this work, the author used a novel bioinformatics tool to find possible SNP in the amino acid sequence of AKAP9 and found no SNP. This might imply that the AKAP9 might have no polymorphism. However, due to the limitation of the present tool, the conclusion cannot be reached. Future analysis by future available tool is suggested. In addition, a study on the probable mutated prone position, weak linkage analysis, is another *in silico* methods suitable for this AKAP protein.

Declaration on Conflict of Interest

The author hereby would like to declare for no conflict of interest in this work.

References

- Borgiani P, Ciccacci C, Forte V, Romano S, Federici G, Novelli G (2007) Allelic variants in the CYP2C9 and VKORC1 loci and interindividual variability in the anti-coagulant dose effect of warfarin in Italians. *Pharmacogenomics* 8: 1545-50.
- Burns HLL, Barraclough DM, Taylor SS (2004) Identification and functional analysis of dual-specific A kinase-anchoring protein-2. *Methods. Enzymol* 390: 354-74.
- Chen L, Marquardt ML, Tester DJ, Sampson KJ, Ackerman MJ, Kass RS (2007) Mutation of an A-kinase-anchoring protein causes long-QT syndrome. *Proc Natl Acad Sci USA* 104: 20990-5.
- Chen L, Kass RS (2006) Dual roles of the A kinase-anchoring protein Yotiao in the modulation of a cardiac potassium channel: a passive adaptor versus an active regulator. *Eur J Cell Biol* 85: 623-6.
- Frank B, Wiestler M, Kropp S, Hemminki K, Spurdle AB, Sutter C, Wappenschmidt B, Chen X, Beesley J, Hopper JL; Australian Breast Cancer Family Study Investigators, Meindl A, Kiechle M, Slanger T, Bugert P, Schmutzler RK, Bartram CR, Flesch-Janys D, Mutschelknauss E, Ashton K, Salazar R, Webb E, Hamann U, Brauch H, Justenhoven C, Ko YD, Brüning T, Silva Idos S, Johnson N, Pharoah PP, Pooley KA, Chang-Claude J, Easton DF, Peto J, Houlston R; Gene Environment Interaction and Breast Cancer in Germany Group, Kathleen Cuninghame Foundation Consortium for Research into Familial Breast Cancer Investigators, Australian Ovarian Cancer Study Management Group, Chenevix-Trench G, Fletcher O, Burwinkel B (2008) Association of a common AKAP9 variant with breast cancer risk: a collaborative analysis. *J Natl Cancer Inst* 100: 437-42.
- Gasteiger E, Gattiker A, Hoogland C, Ivanyi I, Appel RD, Bairoch A (2003) ExPASy: The proteomics server for in-depth protein knowledge and analysis. *Nucleic Acids Res* 31: 3784-8.
- Lin JW, Wyszynski M, Madhavan R, Sealock R, Kim JU, Sheng M (1998) Yotiao, a novel protein of neuromuscular junction and brain that interacts with specific splice variants of NMDA receptor subunit NR1. *J Neurosci* 18: 2017-27.
- Mack JT, Brown CB, Tew KD (2008) ABCA2 as a therapeutic target in cancer and nervous system disorders. *Expert. Opin Ther Targets* 12: 491-504.
- Marx SO, Kurokawa J (2006) AKAPs as antiarrhythmic targets? *Handb. Exp Pharmacol* 171: 221-33.
- Rudd MF, Webb EL, Matakidou A, Sellick GS, Williams RD, Bridle H, Eisen T, Houlston RS; GELCAPS Consortium (2006) Variants in the GH-IGF axis confer susceptibility to lung cancer. *Genome Res* 16: 693-701.
- Saucerman JJ, Healy SN, Belik ME, Puglisi JL, McCulloch AD (2004) Proarrhythmic consequences

- of a KCNQ1 AKAP-binding domain mutation: computational models of whole cells and heterogeneous tissue. *Circ Res* 95: 1216-24.
12. Shah SC, Kusiak A (2004) Data mining and genetic algorithm based gene/SNP selection. *Artif Intell Med* 31: 183-96.
13. Tao J, Shumay E, McLaughlin S, Wang HY, Malbon CC (2006) Regulation of AKAP-membrane interactions by calcium. *J Biol Chem* 281: 23932-44.