

14th World Congress on

Infection Prevention and Control

December 06-07, 2018 | Valencia, Spain

Rickettsiae in human and in blood feeding arthropods in Northern Tunisia

Fatma Khrouf¹, Saba Zouari¹, Youma M'Ghirbi¹, Souha Boughatef², Nisaf Ben Alaya², Lamia Ammari³ and Lilia Messadi⁴¹IPT, Tunisia,²ONMNE, Tunisia³LaRabta, Tunisia⁴ENMV, Tunisia

Statement of the Problem: Rickettsioses are considered as emerging infectious diseases. These affections are classically transmitted to humans via arthropods vector bites. In Tunisia, about 200 cases were registered every year. Despite medical importance and longstanding presence of this disease, relationship between Rickettsia species and potential arthropod vectors has been poorly investigated. Based on the epidemiological understood and the control of emerging diseases spread in Mediterranean region, a survey of rickettsial diseases was carried out to have an overview about the circulation of these diseases in Northern Tunisia. Nevertheless, this data is necessary in order to target surveillance and control of this vector-borne disease nationwide.

Methodology & Theoretical Orientation: A total of 2452 ectoparasites (ticks and fleas) infesting domestic animals (dogs, sheep and goats) were collected in five studied cities in which sera and blood of patients, suspected to have rickettsial infection on the basis of clinical criteria, were tested serologically (IFI) and molecularly. Extracted DNA was subject to Rickettsia identification using qPCR. To confirm our findings, some amplified positive samples from vectors and humans were sequenced.

Findings: During our survey (March-November 2015), the seroprevalence of 40 suspected patients was 54%. Three positive biopsies were identified infected by *R. conorii*. The global prevalence of infected ticks and fleas by Rickettsia was 46% and 63% respectively. Specific qPCR showed the infection of 92% of positive fleas by *R. felis*. The molecular sequencing, using 3 target specific genes (*gltA*, *ompA*, *OmpB*), allows the identification of *R. massiliae* in *Rh. turanicus*, *R. helvetica* in *Ixodes ricinus* and *R. aeschlimannii* in *Hyalomma marginatum* and *Hy. excavatum*.

Conclusion & Significance: In terms of public health, this study gives a global vision of the distribution of Rickettsia in human and vectors in endemic regions. Ticks and fleas, both abundant arthropods, seem to be the most significant Arthropoda species carrying Rickettsia agents and may play an important role in maintaining rickettsial infections and their transmission to human. Further investigations in humans and animals are needed to confirm these data.

kfatmamail@yahoo.fr