



Addressing Decision Making on End of Life Care for People with Dementia

Jones K, Birchely G, Clare L, Huxtable R, Walter T and Dixon J

Health and Social Care, WELS, The Open University, Walton Road, Milton Keynes, Buckinghamshire, MK76AA, UK

*Corresponding author: Jones K, Health and Social Care, WELS, The Open University, Walton Road, Milton Keynes, Buckinghamshire, MK76AA, UK, Tel: 07891868695; E-mail: kerry.jones@open.ac.uk

Received date: May 19, 2017; Accepted date: May 24, 2017; Published date: May 31, 2017

Copyright: © 2017 Jones K, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Citation: Jones K, Birchely G, Clare L, Huxtable R, Walter T, et al. (2017) Addressing Decision Making on End of Life Care for People with Dementia. J Comm Pub Health Nurs 3: 178. doi:10.4172/2471-9846.1000178

Abstract

Dementia affects approximately 36 million people worldwide and the number living with dementia is expected to increase to 66 million by 2030 [1]. Dementia is recognised as a public health priority which urgently needs attention by the World Health Organisation [2]. Despite recent national and international policy initiatives, people with dementia continue to receive invasive treatment such as artificial nutrition and hydration and inadequate care towards the end of their lives [3]. In our scoping review of decision making and advanced care planning for people with dementia, we outline some of the challenges of decision making that people with dementia, their families and health professionals face in talking about a complex and sensitive issue.

Keywords: Dementia; End of life care; Death and dying; Proxy decision making

Introduction

While there has been increasing attention upon dementia and end of life care by policy makers and international governments, there has been less focus on the complex physical and psychological needs of dementia. As a consequence, people with dementia may be subject to interventions that they might not otherwise have wanted (resuscitation, artificial nutrition and hydration) if the condition is not recognised as being terminal. Including people with dementia and their families in planning for end of life care can provide opportunities to review and initiate appropriate palliative care.

The WHO [2] describes Advance Care Planning (ACP) as a discussion about preferences for future care between an individual and a care provider in anticipation that the person's condition will deteriorate. Some of the benefits of having this discussion are that it provides an opportunity to discuss and later, initiate timely palliative care [4]. With dementia, it is suggested that these discussions should take place in the early stages of the disease [5]. However, the topic can be distressing for some people and their families especially when health and social care professionals are challenged by the lack of interpersonal skills or training to facilitate such a sensitive discussion or are reluctant to assume responsibility for discussing ACP [6].

We conducted a scoping review studies which have explored decision making processes associated with people with dementia and their families. Eligible studies had to report on decision making at the end of life, who was involved in this discussion and how the discussion

was initiated (health care professional, the person with dementia, the family member) [7-11].

In this review of the evidence concerning EoLC for people with dementia and the decisions which influence care at the end of life. Our review was guided by three research questions a. Who is involved in the decision-making process? b. How are decisions made concerning EoLC and what is the outcome in facilitating a good death? c. In what way do systemic factors impact upon EoLC?

Twenty five eligible studies reported on planning for end of life care and the decisions made by and for people with dementia. The studies demonstrate that there are a number of unique challenges that need to be addressed to ensure that people with dementia receive appropriate, adequate and effective care towards the end of their life.

References

1. Alzheimer's Disease International (2012) World Alzheimer's Report 2012: Overcoming the stigma of dementia.
2. World Health Organisation (2011) Palliative care for older people: Better practice. Rome, Italy.
3. World Health Organisation (2012) Dementia a public health priority. Switzerland.
4. National Council for Palliative Care (2012) 2012 Inquiry: Improving dementia diagnosis rates in the UK.
5. van der Steen JT (2010) Dying with dementia: What we know after more than a decade of research. J Alzheimers Dis 22: 37-55.
6. Sampson EL, Jones L, Thuné-Boyle IC, Kukkastenvemmas R, King M, et al. (2011) Palliative assessment and advance care planning in severe dementia. An exploratory randomized controlled trial of a complex intervention. Palliat Med 25: 197-209.
7. Allen RS, DeLaine SR, Chaplin WF, Marson DC, Bourgeois MS, et al. (2003) Advance care planning in nursing homes: Correlates of capacity and possession of advance directives. Gerontologist 43: 309-317.
8. Beauchamp TL, Childress JF (2008) Principles of biomedical ethics. Oxford University Press, New York.
9. Dening KH, Jones L, Sampson EL (2012) Preferences for end of life care: A nominal group study of people with dementia and their family carers. Palliat Med 27: 409-417.
10. Kaufman SR (2005) ... and a Time to die. How american hospitals shape the end of life. Scribner, London.
11. van der Steen JT, Radbruch L, Hertogh CM, de Boer ME, Hughes JC, et al. (2013) White paper defining optimal palliative care in older people with dementia: A Delphi study and recommendation from the European association for palliative care. Palliat Med 28: 197-209.