



Alzheimer's Disease Developed Significantly Less Frequently in So-called "Children of Blockade" Leningrad"

Igor Kvetnoy*

Department of Pathology, Ott Institute of Obstetrics, Gynecology and Reproductology, Saint-Petersburg, Russian Federation

Editorial

Alzheimer's disease (AD) is characterized by progressing loss of the memory, bright dementia and is one of the most serious neurodegenerative frustration. Now AD is observed in more than 20 million people all over the world, and their quantity as it is expected, will be doubled in 25-30 years [1,2].

Now two molecules are thought to be the most important in the mechanism of AD development: β -amyloid and tau-protein [3]. Existence of pathogenetic connection between β -amyloid accumulation and neurofibrillial damages was proved in transgenic mice. The selective neuronal destruction was marked in departments of a brain in which the maintenance of amyloid plague was mostly expressed what allows to assume direct participation of β -amyloid in the process of neuronal death in AD in this mice [2].

Tau-protein is the stabilized phosphoprotein, functionally linked with microtubules and providing the structurally functional cytoskeleton organization [4]. In the patients with AD tau-protein, despite to healthy people, phosphorylating, changes the spatial configuration and aggregate into spiral filaments and by that may not support cytoplasmical microtubules formation [3].

Currently, the diagnosis of a specific form of dementia as Alzheimer's disease (AD) manifestation is confirmed finally only at autopsy using specific immunohistochemical methods [2]. Undoubtedly, the problem of life-time diagnosis of AD is one of the most actual problem of biomedicine. This circumstance dictates the necessity to develop life-time markers for diagnosis and prognosis of AD [1,3-5].

We suppose that the study of extra-brain tissues and cells which are available for biopsy could be very promising for this purpose. Most publications on the use of peripheral cells for diagnosis of AD are devoted to study skin fibroblasts. From our point of view the peripheral blood lymphocytes (PBL) could be considered as more suitable samples for the research in this field of study for many reasons (one of them is the ability of lymphocytes to produce biologically active substances, which could be involved in the pathogenesis of AD).

In 2000 we have tried to detect the expression of tau-protein in human peripheral blood lymphocytes (PBL) in patients with AD,

and our attempt was successful: the expression of tau-protein was immunocytochemically shown in PBL in absolute majority of AD patients studied [6]. The International Consortium of AD studying, created in the USA on the basis of National Institute of Ageing and Rejgan's Institute specifies in the recommendations that the pathology of tau-protein is one of the most important components of AD pathogenesis.

During last 10 years we continue and extend the research in this field on the great cohort of patients, and we found a very interesting fact: the AD developed significantly less frequently in the group of so-called "children of blockade" Leningrad" as well as tau expression in their lymphocytes was also less pronounced than in patients of the same age, who did not survive this tragedy. The demonstration of clear difference in localization of tau-protein in blood lymphocytes between healthy and sick people testifies to the fact that tau-protein could be considered as a promising marker and blood lymphocytes as a suitable sample for life-time diagnosis of AD.

The reduction in the incidence of Alzheimer's disease in people who have undergone a blockade of Leningrad indicates an increase in the stability of nerve connections and cognitive functions in the brain of people during life in childhood with limited nutrition and severe environmental conditions.

References

1. Blennow K (2017) A Review of Fluid Biomarkers for Alzheimer's Disease: Moving from CSF to Blood. *Neurol Ther* 6: 15-24.
2. De-Paula VJ, Radanovic M, Diniz BS, Fortenza OV (2012) Alzheimer's Disease. *Subcell Biochem* 65: 329-352.
3. Taliib LL, Joaquim HP, Fortenza OV (2012) Platelet Biomarkers in Alzheimer's Disease. *World J Psychiatry* 2: 95-101.
4. Henry MS, Passmore AP, Todd S, McGuinness B, Craig D, et al. (2013) The Development of Effective Biomarkers for Alzheimer's Disease: A Review. *Int J Geriatr Psychiatry* 28: 331-340.
5. Tarawneh R, Holtzman DM (2010) Biomarkers in Translational Research of Alzheimer's Disease. *Neuropharmacology* 59: 310-22.
6. Kvetnoy IM, Hernandez-Yago J, Kvetnaia TV, Khavinson VK, Malinin VV, et al. (2000) Tau-Protein Expression in Human Blood Lymphocytes: Promising Marker and Suitable Sample for Life-time Diagnosis of Alzheimer's Disease. *Neuro Endocrinol Lett* 21: 313-318.

*Corresponding author: Igor Kvetnoy, Ott Institute of Obstetrics, Gynecology and Reproductology, Saint-Petersburg, Russian Federation, Tel: +7 812 328-94-55; E-mail: igor.kvetnoy@yandex.ru

Received: October 19, 2016; Accepted: October 20, 2016; Published: October 27, 2016

Citation: Kvetnoy I (2017) Alzheimer's Disease Developed Significantly Less Frequently in So-called "Children of Blockade" Leningrad". *J Neuroendocrinol Res* 1: e101.

Copyright: © 2016 Kvetnoy I. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.