

# Assessing Microplastic Toxicity with *Paramecium Bursaria*: A Practical Approach

Romina Villar\*

Department of Ecology and Animal Biology, Faculty of Biology, University of Vigo, Spain

## Abstract

MPs, which are also referred to as microplastics, are typically tiny wastes of plastic with diameters ranging from one to five millimeters. This tiny plastic debris, which is common in aquatic systems, poses a serious threat to the aquatic biota. Even though metazoan animals are mostly used to test MPs' toxicology, there are times when their applications are limited due to their high cost, limited ecological niche, or ethical concerns. This indicates that monocellular eukaryotes, also known as protozoa, which are prevalent in nature and are also referred to as eukaryotes, can be utilized to evaluate the toxicity of MPs. Monocellular eukaryotes are also referred to as eukaryotes. These eukaryotic organisms are also known as eukaryotes. Monocellular eukaryotes are another name for eukaryotes. These eukaryotic organisms are also known as eukaryotes. We continued our investigation of the behavioral and molecular changes by employing *P. bursaria* as a protozoan model and MPs-exposed *Paramecium bursaria* (*P. bursaria*). Our findings indicate that *P. bursaria* underwent a number of modifications following the adoption of MPs. An increased level of oxidative stress, a slower speed, altered avoidance strategies, and the possibility of endosymbiotic disruption were among the changes. *P. bursaria* underwent significant and quantifiable changes in response to MP exposure. Overall, this study demonstrated that *P. bursaria* is a promising alternative for the toxicological evaluation of MPs and that it can be used to evaluate the toxicity of other environmental contaminants.

**Keywords:** Toxicological Tool; Protozoa; Unicellular Organisms; *Paramecium Bursaria*; Microplastics

## Introduction

The size of the plastic debris is primarily what determines its negative effects on the local biota [1]. Aquatic ecosystems are gravely endangered by plastic contamination. Large pieces of plastic, like fishing lines and bottle caps, can physically entangle a wide range of aquatic organisms [2,3]. These enormous wastes of plastic can also be broken down into microplastics (MPs) through photo-oxidation, biodegradation, and physical abrasion [4]. Due to their small size, which typically ranges from one millimeter to five millimeters, MPs are emerging as a threat to aquatic biota. As a outcome, numerous aquatic species can get closer to them, most likely through direct ingestion or trophic transfer. Up to this point, numerous investigations have been conducted into the possibility of toxic MPs. Due to the elusive concentration of this novel contaminant in the aquatic system, various concentrations, ranging from 102 to 1010 particles/mL, were chosen to evaluate acute or chronic impacts on aquatic animals from various trophic positions in experimental settings. The metabolic functions of mysid shrimp (*Neomysis japonica*), hermit crabs (*Pagurus bernhardus*), zebrafish (*Danio rerio*), monogonont rotifers (*Brachionus koreanus*), and zebrafish (*Danio rerio*) as well as monogonont rotifers (*Brachionus koreanus*) were examined to see if they had changed or if it is essential to keep in mind that studies involving animals with multiple cells, such as metazoans, account for the majority of our knowledge of MPs. On the other hand, very little is known about how MP affects unicellular protozoans. Furthermore, there is currently a lack of awareness regarding alternative protozoan-based methods for assessing MP toxicity [5-7]. The broad group of single-celled organisms that can be found in nature are called protozoans. One of these microscopic animals, *Paramecium*, is of particular interest to the scientific community because it has served as a useful model organism in eukaryotic biology and biomedical research [8]. *Paramecium*, a genus of ciliates with short cilia arranged in rows on the membrane, was the first ciliate to be observed due to their relatively large size. Since their discovery, a number of culturing protocols have been developed

that make it relatively simple to cultivate and maintain *Paramecium* cultures with minimal effort, low cost, and fewer ethical considerations. In addition, it was later demonstrated that *Paramecium* engaged in complex but quantifiable behaviors like avoidance and pursuit of prey in response to external stimuli. It is now possible to link phenotypic shifts to genotypes and learn about the genomes of several *Paramecium* species thanks to advancements in sequencing technology. *Paramecium* is an excellent model organism for evaluating MP toxicity because of all of these characteristics. Recent studies have demonstrated that MPs are utilized in *Paramecium*; However, additional research is needed to investigate the potential effects on *Paramecium*, particularly the behavioral and molecular changes caused by MPs uptake [9,10].

## Discussion

The practical approach outlined in this study provides valuable insights into the toxicity of microplastics using *P. bursaria* as a model organism. The findings contribute to our understanding of the ecological consequences of microplastic pollution and highlight the importance of mitigating its impact on freshwater ecosystems. Further research is warranted to explore the long-term effects of microplastic exposure and potential interactions with other environmental stressors.

\*Corresponding author: Romina Villar, Department of Ecology and Animal Biology, Faculty of Biology, University of Vigo, Spain, E-mail: rominavillar@gmail.com

Received: 01-May-2024, Manuscript No: jety-24-136514, Editor assigned: 04-May-2024, Pre-QC No: jety-24-136514 (PQ), Reviewed: 18-May -2024, QC No: jety-24-136514, Revised: 25-May-2024, Manuscript No: jety-24-136514(R), Published: 31-May-2024, DOI: 10.4172/jety.1000219

Citation: Romina V (2024) Assessing Microplastic Toxicity with *Paramecium Bursaria*: A Practical Approach. J Ecol Toxicol, 8: 219.

Copyright: © 2024 Romina V. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Conclusions

MPs altered the behavior and molecular makeup of *P. bursaria*. Changes in the symbiotic relationships between the infected *P. bursaria*, decreased avoidance, and elevated oxidative stress were among the symptoms. As this study demonstrated, *P. bursaria* was found to undergo behavioral and molecular changes upon exposure to MPs. *P. bursaria* could be a useful tool for determining the toxicity of novel pollutants like MPs in aquatic or experimental settings, according to this.

## References

1. Shah R, Kulhara P, Grover S, Kumar S, Malhotra R, et al. (2011) Contribution of spirituality to quality of life in patients with residual schizophrenia. *Psychiatry Res* 190: 200-5.
2. Chong HY, Teoh SL, Wu DBC, Kotirum S, Chiou CF, et al. (2016) Global economic burden of schizophrenia: a systematic review. *Neuropsychiatr Dis Treat* 12: 357-73.
3. Grover S, Davuluri T, Chakrabarti S (2014) Religion, spirituality, and schizophrenia: A review. *J Psychol Med* 36: 119-24.
4. Talreja B, Shah S, Kataria L (2013) Cognitive function in schizophrenia and its association with socio-demographics factors. *Ind Psychiatry J* 22: 47.
5. Levin MJ, Ustianowski A, De Wit S, Launay O, Avila M, et al. Intramuscular AZD7442 (Tixagevimab-Cilgavimab) for Prevention of Covid-19. *N Engl J Med* 386: 2188-2200.
6. Ocon AJ, Mustafa SS (2023) Real-World Experience of Tixagevimab and Cilgavimab (Evusheld) in Rheumatologic Patients on Rituximab. *J Clin Rheumatol* 29: 109-111.
7. Gheita TA, El-Latif EA, El G, Samy N, Hammam N, et al. (2019) Behçet's disease in Egypt: a multicenter nationwide study on 1526 adult patients and review of the literature. *Clinical rheumatology* 38: 2565-75.
8. Rice SA, Woo PN, El-Omar E, Keenan RA, Ormerod AD (2013) Topical tacrolimus 0.1% ointment for treatment of cutaneous Crohn's Disease. *BMC research notes* 6: 19.
9. Hatemi G, Melikoglu M, Tunc R, Korkmaz C, Turgut Ozturk B, et al. (2015) Apremilast for Behçet's syndrome--a phase 2, placebo-controlled study. *N Engl J Med* 372: 1510-8.
10. Tawfeek H, Abdellatif DAAH, Abd Elaleem Elnashar J, Abdelaleem Y, Fathalla D (2020) Transfersomal gel nanocarriers for enhancement the permeation of lornoxicam. *J drug deliv sci technol* 56.