

Assessing of Knowledge, Attitude and Practice of Sunlight Exposure of Infants among Mothers Attending in Nigest Elleni Mohammed Memorial Comprehensive Specialized Hospital, Hossana Ethiopia 2023

Eneyew Gedamu* and Awgchew

Department of Pediatric Nursing, Wachemo University, Ethiopia

Abstract

Background: Sun exposure has the most important benefit: it increases the amount of vitamin D absorbed into your body, also sunlight exposure is the main source of vitamin D. Daily exposure to sunlight remains the cheapest, safest, and most effective method for preventing vitamin-D-related diseases.

Objective: To Assessing of knowledge, Attitude, and Practice of Sunlight Exposure of Infants among Mothers Attending in Nigest Elleni Mohammed Memorial Comprehensive Specialized Hospital Hosanna Town Ethiopia 2023.

Method: Institutional base cross-sectional study was conducted among mothers of infants visiting NEMMCSH from November1-December30/2023.to select the study participant systematic random sampling was applied. Data were entered and analysed by using SPSS version 24.

Result: A total of 252 study participants were served with a response rate of 100% of these 107 (42.4%) of them aged between 25-29yrs, 171(67. 9%) of the mothers could read and write, similarly 196 (77. 9%) of their husbands could also read and write. The majority of the respondents 238 (94.4%) knew about sunlight exposure of their infants. Among 252 respondents 232 (92.1%) of them knew of morning is a good time to expose to sunlight. Almost all (99.6%) of mothers had a positive attitude towards sunlight exposure. Of the total respondents (90.1%) of mothers believed that sunlight exposure helps to prevent rickets in infants. Most of the respondents 227(90.07%) exposed directly or indirectly of their infants to sunlight. 165 respondents started to expose their child to sunlight between 16 -30 days and More than half of participants (70%) exposed their Children to sunlight

Conclusion: According to our study, the majority of the respondents (95%) had good knowledge about sunlight exposure of their infants and 98% of respondents had good attitudes. On the other hand, 29.7 % of mothers disagree about the benefit of sunlight exposure to their infant, from these, 49.2% of mothers believe sunlight exposure to infants can cause evil eye.

Keywords: Sunlight; Exposure; Knowledge; NEMMCSH

Introduction

Regular exposure to sunlight increases serotonin levels in the body, making the body more alert. Sun exposure has the most important benefit: it increases the amount of vitamin D absorbed into your body, also sunlight exposure is the main source of vitamin D [1]. Daily exposure to sunlight remains the cheapest, safest, and most effective method for preventing vitamin-D-related diseases. Exposing baby to sunlight early in the morning is a good way to prevent rickets and reduce the problem of vitamin D deficiency and jaundice in newborns .

Sunlight prevents and cures rickets by discovering an inactive lipid in food and skin that is converted into anti-rickets by UV light. The benefits of good sun exposure are primarily increased vitamin D introduced into the body. Most cases of V-D deficiency are caused by inadequate exposure to sunlight, the active form of the vitamin, some of which are involved in calcium metabolism and neuromuscular activity of the immune system. On the other hand, Vitamin D deficiency results in growth retardation and skeletal deformities in children, infantile eczema in neonates, and may result in muscle weakness and bone fractures in adults. Rickets are unacceptably high worldwide, with the greatest burden in Africa, the Middle East, and Asia. An estimated one billion people worldwide, across all age groups, have a vitamin D deficiency and almost 50% of the world population lives with Vitamin D insufficiency. Disease related to less awareness of proper sunlight exposure and low knowledge, attitude & practice towards it, is becoming a significant burden to developing children,

especially among breastfeeding infants [2].

In addition, epidemiological studies have revealed that vitamin-D deficiency is highly prevalent among infants in different countries, regardless of age, ethnicity, geographical location, and climatic conditions, and is believed to range from 2.7% to 45% mostly due to inadequate practice of mothers exposed to sunlight which is common mainly in developing countries especially in Sub-Saharan African. World Health Organization report estimates that globally one billion people have vitamin-D deficiency and this burden also increases major disorders of the musculoskeletal system and possibly an increased risk of various autoimmune diseases and cancers.

In addition, vitamin D deficiency has also a substantial economic burden as much as 25% of health care dollars could be saved just by

*Corresponding author: Eneyew Gedamu, Department of Pediatric Nursing, Wachemo University, Ethiopia, Email: ewseife@gmail.com

Received: 01-Jan-2024, Manuscript No. jpch-24-127872; **Editor assigned:** 03-Jan-2024, PreQC No. jpch-24-127872 (PQ); **Reviewed:** 17-Jan-2024, QC No. jpch-24-127872; **Revised:** 23-Jan-2024, Manuscript No. jpch-24-127872 (R); **Published:** 31-Jan-2024, DOI: 10.4172/2376-127X.1000618

Citation: Gedamu E (2024) Assessing of Knowledge, Attitude and Practice of Sunlight Exposure of Infants among Mothers Attending in Nigest Elleni Mohammed Memorial Comprehensive Specialized Hospital, Hossana Ethiopia 2023. J Preg Child Health 11: 618.

Copyright: © 2024 Gedamu E. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

improving the world's vitamin D status.

Recent studies have shown that vitamin D insufficiency is common in tropical countries even if there is sunshine throughout the year. Even though a high proportion of sunny weather in Ethiopia the occurrence of rickets is very highly which is 40% is due to mal-practice and inadequate knowledge towards infant's sunlight exposure and due to broad studies were not conducted to assess the knowledge of mothers on adequate sunlight exposure which contributes to infant mortality and morbidity [3]. Ethiopia adopted health education as a strategy to change maternal behavior on infants' sunlight exposure to prevent rickets in the country during the 1960s. Implementation of this strategy remains low and inconsistent, and health messages had no focus that positively influences maternal practice and infants not get adequate sunlight. Lack of sufficient information on the determinant of this risky behavior and the attitude of Ethiopian mothers initiated this study focus on assessing of knowledge, attitude, and practice of sunlight exposure of infants among mothers attending in NEMMCSH in Hosanna town.

Methods

Study area and period

The study was conducted in Nigist Eleni Mohamed Memorial Compressive Specialized Hospital from November 1-december 30/2023. The hospital is found in Hosanna town of Hadiya zone. It is a teaching hospital which provides service for more than 1.5 million catchment populations and a total capacity of 250 beds. Also it has 590 health professionals [4].

Study Design

Institution based cross sectional study was conducted.

Population

Source population

All women whose child's age was less than 1 year and visit NEMMCSH

Study population

All women whose child's age were less than 1 year and visiting NEMMCSH for health care and immunization during the study period.

Sampling procedure

Systematic random sampling method was applied to identify study participants. To get study participants, first the average number of lactating mothers who were visit NEMMCSH was 780, which was obtained by referring a two-month registration report. The k^{th} interval was determined by dividing the total population size by the total sample size. $K = N/n$; $K = 780/252 = 3$.

Study variables

Independent variables

Knowledge, attitude, and practice of sunlight exposure of infant.

Dependent variable

Age, Sex, Religion, Cultural, Educational status, Occupation, Marital status, sex of infant, and age of infant [5].

Operational definitions

Knowledgeable/good knowledge: Mothers who got above the median score from the knowledge question were considered as knowledgeable.

Good practice: Participants were deemed to have had good practice if they received a score on the practice questions of greater than or equal to the median.

Poor practice: Mothers scored less than median of the total score of observational checklist criteria.

Good attitude: Mothers who got above the median score from the attitude question were considered as good attitude.

Poor attitude: Mothers who got below the median score from the attitude question were considered as poor attitude.

Data collection Instrument and procedure

Data was collected through face-to-face interviews after preparing a structured and pre-tested questionnaire. The questionnaire was initially prepared in English and translated into Amharic and again retranslated back to English to check for any inconsistencies or distortions in the meaning of words and concepts [6-8].

Data quality control

To ensure the data was maintained through a common understanding of the tool among group members, a pretest was checked for its completeness before actual data collection, Processing, and analysis. Two-day intensive training about tool utilization, ethics, data collection methods, and the study's purpose was given to data collectors and supervisors.

Discussion

This study was focused on assessing the knowledge, attitude and practice among mothers of infants in NEMMCSH. In the study 94.4% of mothers of infants had information (knowledge) about infant sunlight exposure; the same study done in England showed that 98.6% of the mothers had information about sunlight exposure. So the result of our study is lower than the study done in England the possible reason may be due to educational differences in the study participants and economic and developmental differences between the countries. Also differences in the use of technology. In our study, 96.8% of the respondents mentioned that "sunlight exposure had benefits this finding was higher than the study done in Farta district of the south Gonder zone. Which was 75.98% the difference may be due to our study's 94.4% of the mothers had information (knowledge) about sunlight exposure. On the other hand, our result is lower than the study done in Jimma Town.

The percentage of respondents who reported that sunlight exposure was beneficial for strengthening bone was 53.5% this finding was in line with a study done in Jimma and Turkey. Regarding the best time for sunlight exposure of infants, 92.1% of the mothers mentioned it was in the morning; also 6.3% of them mentioned in the afternoon and the remaining 1.6% of the respondent in the evening this was similar to the same study done in Aleta wondo. Which was 90.4% this similarity may be due to the socio-cultural similarity between the study sites and both study areas were found in the same region. On the other hand, the result was higher than the study done in Farta district. The study findings show that 99.2% of the respondents had a positive attitude to infant sunlight exposure the result was higher than the study done in Debre Tabor Town which was 61% the reason may be due to difference in study population and study area and also feeling of respondents

while they exposed their infants to sunlight 89.7% of them were happy and 10.3% of the mother were had anxiety which is higher than study done in Ethio-Swedish hospital and Farta district the possible reason may be due to socio-natural difference.

The current study identifies 14% of the mothers perceived sunlight exposure had a bad effect on infants from these 8.6% of them think had a respiratory problem, 14.3% of them thought it was exposure to the evil eye and 5.7% of them thought it may cause a mental problem this result lower than a similar study conducted in Debre-Tabor Twon. Regarding the practice of sunlight, 99.6% of the mothers exposed their infant to sunlight this finding was in line with the study done in Dessie Town which was 93.9% but higher than the study done in Yergalem(84%) respectively. In this study 65.5% of the respondents started to expose their infant to sunlight between 16-30 days the result was higher than the study done in Debre-Berehan, Dessie and Yergalem which was 12.6%, 19.3% and 25.6% respectively. This discrepancy may be due to participants source of information in our study 76.5% of respondents were got information from health professionals [9].

In our study, 96.4% of the respondents exposed their infant to sunlight in the morning before 4 am this result was in line with studies conducted in Debre-Markos, Debre-Berhan and Dessie which was 99.1%, 97% and 93.4% respectively. This may be due to near similar topography of the study area.

The findings of this study show that 83.3% of the mothers unclosed their infants when they were exposed to sunlight the results from Metu oromia, Debre-markose, Debre-Berhan and Dessie were lower than when we compared with this finding. On the other hand, 95.2% of the respondents applied lubricant or lotion during sunning their infant, the result was in line with a study done in Metu Oromia. This implies that it is crucial to increase the awareness of communities regarding the appropriate practice of sunning infants for optimal cutaneous vitamin D production.

Limitation of the study

Owing to a lack of funding and resources, all medical facilities were no included to the study, which could have decreased the study's accuracy.

The mother did not exhibit any conduct, and the information was self-reported.

Insufficient funds and time were available to complete this study.

Owing to the nature of descriptive research, qualitative study design is preferable [10-12].

Conclusion

According to our study, the majority of the respondents (95%) had good knowledge about sunlight exposure of their infants and 98% of respondents had good attitudes. On the other hand, 29.7 % of mothers disagree about the benefit of sunlight exposure to their infant, from these, 49.2% of mothers believe sunlight exposure to infants can cause evil eye. Therefore, health education focusing on the importance of sunlight exposure is important to improve knowledge, practice, and attitude of mothers' sunlight exposure of their infants.

Ethical Consideration

Ethical clearance for the commencement of the study was obtained from Wachemo University, research and community service vice

president's office Research Ethical Committee, Permission was also sought and obtained from the ethical committee of Nigest Eleni Mohammed Memorial specialized hospital. Data was kept anonymous by keeping the identity of the neonate's or mother's credentials hidden before, during, and after the study.

Funding: This research didn't receive any grant from any funding agency in the public, commercial or not-for-profit sectors

Authors' contributions

All authors collaboratively made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

Abbreviations

NEMMCSH: Nigest Eleni Mohammed memorial comprehensive specialized hospital.

SLE: Sun light exposer.

UV- light: Ultraviolet light.

Conflicting interests

No potential conflict of interest

References

1. Minari JB, Okeke U (2014) Chemopreventive effect of *annona muricata* on DMBA-induced cell proliferation in the breast tissues of female albino mice. *Egy J Med Human Genet* 15: 327.
2. Baskaran N, Manoharan S, Balakrishnan S, Pugalendhi P (2010) Chemopreventive potential of ferulic acid in 7,12-dimethylbenzaanthracene-Induced mammary carcinogenesis in Sprague-Dawley rats. *Eur J Pharmacol* 63: 22.
3. Sharmila R, Manoharan S (2012) Anti-tumor activity of rosmarinic acid in 7, 12-dimethylbenz (a) anthracene (DMBA) induced skin carcinogenesis in Swiss albino mice. *Ind J of Physio Sciences* 7: 344-356.
4. Suresh S, Manoharan M, Vijayaanand P, Sugunadevi A (2020) Chemopreventive and antioxidant efficacy of (6)-paradol in 7, 12-dimethylbenz (a) anthracene induced hamster buccal pouch carcinogenesis. *Pharmacological Reports* 62: 1178-1185.
5. Anjugam C, Sridevi N, Rajendra Prasad M, Balupillai A (2018) Morin prevents ultraviolet-b radiation-induced photocarcinogenesis through activating thrombospondin-1 in the mouse skin. *Asian J Pharma Clin Res* 11: 24-34.
6. Farhan M, Khan M, Oves H (2016) Cancer therapy by catechin involves redox cycling of copper ions and generation of reactive oxygen species. *Toxins* 8: 37.
7. Li B, Zhao J, Wang C (2011) Ginsenoside Rh2 induces apoptosis and paraptosis-like cell death in colorectal cancer cells through activation of P53. *Cancer Letters* 301: 185.
8. Ahire V, Kumar K, Mishra, Kulkarni P (2017) Ellagic acid enhances apoptotic sensitivity of breast cancer cells to c-radiation. *Nutri & Cancer* 69: 904-910.
9. Liang F, Li A, Shi R, Liu J, Tang Y, et al. (2018) Anticancer effects of ginsenoside Rg1, cinnamic acid, and tanshinone IIA in osteosarcoma MG-63 cells: nuclear matrix down regulation and cytoplasmic trafficking of nucleophosmin. *Inte J Bioche & Cell* 5: 567-574.
10. Cowling VH, Cole MD (2017) E-cadherin repression contributes to c-Myc-induced epithelial cell transformation. *Oncogene* 26: 3582-3586.
11. Clark GM (2015) Prognostic and predictive factors for breast cancer. *Breast Cancer* 2: 79-89.
12. Ariazi EA, Ariazi JL, Cordera F, Jordan VC (2016) Estrogen receptors as therapeutic targets in breast cancer. *Curr Medicinal Che* 6: 181- 202.