



## Detection of P53 Gene of Colorectal Cancer and Activating it by Stabilizing Dynamic Cerna

Avery Nelson\*

Department of Cardiology, University Hospital Heidelberg, Germany

### Introduction

Colorectal cancer (CRC) is one of the deadly cancers in the world. It stands in the third place with 1,933,600 numbers of patients according to 2020 analysis. There are other names for CRC like colon cancer, bowel cancer, rectal cancer or adenocarcinoma because they emerge from the glandular epithelial cells of large intestine at the colon and rectum.

A person may be affected with CRC due to following reasons

- Family history of CRC
- Genetic disorders
- Certain health conditions like Inflammatory bowel disease
- Type 2 Diabetes

CRC gets spread to some other parts of body by metastasis once the condition becomes severe and then no further medication can help the patient.

A tumour suppressor gene (TP53) is usually involved in control of apoptosis and cell cycle. This is located on the short arm of chromosome 17 (17p13.1) through this communication it is explained that how TP53 gene is targeted by using chemically stabilised endogenous RNA.

Competitive endogenous RNA is a previously considered as junk but after several studies it is discovered that they code for lincRNA. ceRNA is a network of several mRNAs which competes for same pool of micro RNAs. To regulate the ability of miRNA and to inhibit the translation of mRNA into proteins the ceRNA binds to their target mRNAs and acts as miRNAs sponges. This is called as sponging mechanism.

The action site of p53 gene in CRC is first identified and then miRNAs will block translation by binding to mRNA. By using chemically stabilised ceRNA the excess mi RNAs are sponged. Since the ceRNA is a complex secondary structure with many loops and stems it is impossible to perform sponging mechanism by itself sometimes. Hence, two strategies are performed to make them able to perform sponging mechanism.

### Designing ceRNA by chemical Biology

A ceRNA has many nucleotides since it is a complex and dynamic structure. In this method the covalent bonds between all the nucleotides are made stronger. The very first step involves the synthesis of ceRNA by designing artificial nucleotides in the stem of the loop. Later the bonds will be locked by using a light source at a particular wavelength so that they will stay intact as per requirement and do not change. As a result strong covalent bonds are formed.

### Click chemistry reaction

Azide-alkyne cyclo addition which contains triazoles is used to design ceRNA by click chemistry reaction. It is stereospecific and region selective 5 membered ring structure. To stop the conformational changes in dynamic structure of ceRNA, we will use this heterocyclic ring structure of azide alkyne and lock the stem region at our required

site on the ceRNA

### Conclusion

The R1 and R2 groups on the heterocyclic compound will bind at specific bases and interlock them. After this process, by using synthetic biology a cell is designed with help of genetic circuits to generate the ceRNA at the TP53 gene site and activate it.

### Acknowledgement

I would like to thank my Professor for his support and encouragement.

### Conflict of Interest

The authors declare that they are no conflict of interest.

### References

1. Ewing I, Hurley JJ, Josephides E, Millar A. (2014) The molecular genetics of colorectal cancer. *Frontline Gastroenterology* 5:26–30.
2. Vogelstein B, Fearon ER, Hamilton SR. (1988) Genetic alterations during colorectal-tumour development. *N Engl J Med* 319:525–32.
3. August DA, Ottow RT, Sugarbaker PH. (1984) Clinical perspective of human colorectal cancer metastasis. *Cancer Metast Rev* 3: 303–324.
4. Ergun S, Oztuzcu S, Oncocers. (2015) CeRNA-mediated cross-talk by sponging miRNAs in oncogenic pathways. *Tumor Biol* 36: 3129–3136.
5. Rawla P, Sunkara T, Barsouk A. (2019) Epidemiology of colorectal cancer: incidence, mortality, survival, and risk factors. *Gastroenterology Review/Przegląd Gastro enterologiczny* 14:89-103.

\*Corresponding author: Avery Nelson, Department of Cardiology, University Hospital Heidelberg, Germany, Tel: +41 2302665972; E-mail: anelson@med.uni-heidelberg

Received: 18-Oct -2022, Manuscript No. jbtbm-21-44790; Editor assigned: 20-Oct -2022, PreQC No. jbtbm-21-44790 (PQ); Reviewed: 12-Jan -2022, QC No. jbtbm-22-50645; Revised: 19-Jan-2022, Manuscript No. jbtbm-21-44790 (R); Published: 24-Jan-2022, DOI: 10.4172/2155-952X.1000255

Citation: Nelson A (2022) Detection of P53 Gene of Colorectal Cancer and Activating it by Stabilizing Dynamic Cerna. *J Biotechnol Biomater*, 12: 255.

Copyright: © 2022 Nelson A. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.