

Epidemiological Profile of Mental Health Program Patients in Amazon Health Unit

Amanda MR¹, Elisa MNB¹, Juliana MMDO¹, Francisco ITRJ², Andrew MM¹ and Vanessa NB^{3*}

¹Department of Epidemiology, State University of Para, Para

²Department of Medicine, Federal University of Para, Para

³Department of Epidemiology, Federal University of Para, Para

Abstract

Aim: To verify the epidemiological profile of patients enrolled in the Mental Health Program at a Family Health Unit, as well as the most prevalent disorders, age and comorbidities.

Methods: After analyzing the medical records, the following variables were obtained: diagnosis of mental health status, gender, age group, associated comorbidities and medication in use.

Results: There was a prevalence of 61.3% in women and 38.7% in men; patients aged 30 to 39 years were the most affected (22.6%); the most common disorders were epilepsy (36%), depression (14%), anxiety disorder (8.6%) and schizophrenia (8.6%); the most common comorbidities were systemic arterial hypertension (30%) and insomnia (15%) and the most commonly used psychotropic drugs were amitriptyline, carbamazepine and phenobarbital (10.2% each).

Conclusion: There was a high prevalence of associated comorbidities and greater involvement of men aged 30 to 39 years.

Keywords: Mental health; Mental disorders; Comorbidity; Epidemiology; Prevalence, Primary health care

Introduction

Since the 1990s, epidemiology has begun to highlight new demands for care such as mental health, as mental health problems have quietly become the leading cause of disability, morbidity and premature death, in both developed and developing countries [1].

Countries all around the world are going through this "epidemiological transition". Diseases such as depressive disorders and cardiovascular disorders are rapidly replacing the prevalence of malnutrition and infectious diseases. However, this process is occurring without the proper awareness of experts in service planning and public health care. Therefore, mental health care is not exempt from the precariousness of humanized treatment [2].

Despite the high prevalence, myths about mental illness and stigmatization persists even among some healthcare professionals who are unaware of the great progress that has been made in recent decades regarding the diagnosis and, especially, treatment of these psychiatric disorders. About the interventions against the stigma of mental illness, one of the major positive impact factors is prior contact by the intervention participant with someone with mental illness [3,4].

The lack of information about Mental Health actions in Primary Care is evident. In contrast, the patient with mental disorders is guaranteed by law No. 10.216 and has the right to access the best treatment that health system can provide, concerning their needs, and to be treated with humanity and respect with sole interest of benefiting their health, aiming to achieve their recovery through insertion in family, work and community [5].

Since Primary Care is the gateway to Brazilian public health system, the insertion of Mental Health actions allows psychic symptoms to be detected and treated early, avoiding unnecessary hospitalizations [6]. In this sense, the Family Health Program (FHP) is the main strategy to enable primary health care, through the proposal of change in the

rationality of care, based on a multiprofessional teamwork process in which health surveillance, comprehensiveness of practices, hierarchization, territorialization and enrollment of those registered in the respective program [7].

In addition, current national mental health policy consists with progressive reduction of psychiatric beds and expansion and strengthening of out-of-hospital network, consisting mainly of the Psychosocial Care Centers (CAPS) and Therapeutic Residential Services (SRT). It also includes the importance of mental health actions in primary care [8]. CAPS is a type of community health service that provides daily care to patients with severe and persistent mental disorders, under intensive, semi-intensive and non-intensive treatment, according to severity of clinical condition [9].

In fact, mental health deserves specific attention, given its increasing prevalence in world population. Therefore, considering the scarcity of information about Mental Health actions in primary care, it is necessary to contribute with data about the clinical / epidemiological profile of psychiatric patients treated at the Primary Care of Brazilian Unified Health System (SUS). Thus, the objective of this study is to verify the epidemiological profile of patients enrolled in the Mental Health Program in a FHU, as well as identify the most prevalent disorders, age and comorbidities.

*Corresponding author: Vanessa Novaes Barros, Department of Epidemiology, Federal University of Para, Para, Tel: 559199265747; E-mail: barros.novaes@hotmail.com

Received November 06, 2019; Accepted July 15, 2020; Published July 22, 2020

Citation: Amanda MR, Elisa MNB, Juliana MMDO, Francisco ITRJ, Andrew MM, et al. (2020) Epidemiological Profile of Mental Health Program Patients in Amazon Health Unit. *Epidemiol Sci* 10: 379.

Copyright: © 2020 Amanda MR, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Methods

The patients of this research were studied according to the precepts of the Declaration of Helsinki and the Nuremberg Code, respecting the Research Standards Involving Human Beings (Res. 466/12) of the National Health Council, after approval by the direction of the Family Health Unit (USF), in addition to the commitment made by the researchers about the confidentiality of the data used, based on the Term of Commitment to Data usage.

This research is characterized as a cross-sectional, observational, descriptive study, which aimed to evaluate patients enrolled in the USF Mental Health Program. The study included patients of both sexes, without age restriction, without distinction about race, educational level, income levels or marital status. These patients should be registered with USF and the Mental Health Program by November 2015.

The study excluded patients whose data in the medical records were not correctly completed and accessible to researchers, or those who were outside the established period for analysis.

The research was developed from the analysis of medical records at the Health Unit, from November to December 2018. This analysis included research on epidemiological variables: diagnosis of mental health status, gender, age group, associated comorbidities and medication. in use according to the research protocol model elaborated. The data was organized in tables using Microsoft Excel software and was submitted to a descriptive analysis.

Results and Discussion

In this study, was observed a higher prevalence among women as shown in Table 1. These data disagree with previous findings of a survey conducted with 97 medical records of a psychiatric unit in the city of Jequié, Bahia, where 54.6% of patients were actually male [10].

According to Rennó Jr. et al. from menarche to post-menopause, some women suffer from specific disorders, including premenstrual dysphoric disorder, perinatal and perimenopausal depression, blues and puerperal psychosis, as well as mood and anxiety disorders associated with infertility and aborted pregnancies, and also present higher incidence of anxiety, eating disorders, autoimmune diseases and pain [11].

In this research, we also identified higher prevalence of disorders such as depression and generalized anxiety in females as shown in Tables 2 and 3. This can be explained by female tendency to internalize stress in certain situations such as family instability, which leads to disappointment, anxiety, insecurity and affective deficits, culminating in psychological imbalance those results in depression [11].

Women are also frequently exposed to stressors such as violence, abuse and rape since an early age, but not all women who encounter such situations develop disorders. Thus, the interaction between genetic vulnerability, environmental, physiological and psychosocial factors, neurotransmitter and neuroendocrine functioning plays an important role in their pathogenesis [11].

Gender	N	%
Female	38	61.3
Male	24	38.7
Total	62	100

Source: Research protocol

Table 1: Gender of patients registered in the USF Mental Health Program in 2018 Ananindeua - PA.

Diagnosis	N	%
Epilepsy	25	36
Depression	10	14
Anxiety disorder	6	8.6
Schizophrenia	6	8.6
Panic Syndrome	3	4.3
Cerebral Palsy	3	4.3
Behavior disorder	2	2.9
Psychosis	2	2.9
Aggressive Disorder	1	1.4
Obsessive-compulsive disorder	1	1.4
Bipolarity	1	1.4
Alzheimer's disease	1	1.4
Psychomotor agitation	1	1.4
Autism	1	1.4
Parkinson's disease	1	1.4
Not specified	6	8.6
Total	70	100

Source: Research protocol

Table 2: Diagnosis of mental disorder of patients enrolled in the USF Mental Health Program in 2018. Ananindeua-PA*

*Each patient may have more than one diagnosis

Medication	N	%
Amitriptyline	12	10.2
Carbamazepine	12	10.2
Phenobarbital	12	10.2
Risperidone	9	7.7
Haloperidol	9	7.7
Diazepam	8	6.85
Biperiden	8	6.85
Levomepromazine	6	5.1
Clonazepam	5	4.3
Phenytoin	5	4.3
Promethazine	5	4.3
Paroxetine	4	3.4
Chlorpromazine	4	3.4
Valproic acid	4	3.4
Pipothiazine Palmitate	2	1.71
Sertraline	2	1.71
Topiramate	1	0.85
Gabapentin	1	0.85
Memantine Hydrochloride	1	0.85
Oxcarbazepine	1	0.85
Clobazam	1	0.85
Escitalopram	1	0.85
Periciazine	1	0.85
Thioridazine	1	0.85
Donepezil Chloridate	1	0.85
Discontinued Treatment	1	0.85
Total	117	100

Source: Research protocol

Table 3: Psychoactive medications used by patients enrolled in the USF Mental Health Program in 2018. Ananindeua-PA*

*Each patient may use more than one medication.

The association between psychiatric disorders and metabolic disorders has long been described. Individuals with mental disorders constitute a group of high morbidity and mortality. The standardized mortality rate of individuals with schizophrenia is 1.5 to 4 times the general population, and deaths from natural causes account for almost two thirds of this excess, with a significant increase in mortality indicators related to diabetes and cerebrovascular disease. Depression

confers a 1.5 to 2.5 fold relative risk for morbidity and mortality from cardiac causes. It is postulated that the serotonergic system is involved in the association between metabolic syndrome and depression. Decreased serotonergic function is associated with increased carbohydrate intake, bulimic episodes and obesity [12].

To relieve such depression as well as symptoms of other mental disorders, it is common for patients to adopt a high calorie diet (with excessive carbohydrates and fats and low in fiber) due to its anxiolytic effect caused by reduction of hormone levels [13]. However, such practice often leads to excessive weight gain, which in turn culminates in other clinical complications such as systemic arterial hypertension, one of the most prevalent among these patients [14], as evidenced in Figure 1. In addition, many psychiatric drugs have important side effects related to metabolic syndrome, such as weight gain, changes in glucose metabolism and dyslipidemia [12].

The existence of an association between severe mental disorders and high blood pressure is questionable, particularly in patients with schizophrenia. However, the presence of depressive symptoms may be a predisposing factor for the onset of hypertension. Two prospective studies specifically assessed this topic and found increased risk of hypertension in patients with high scores of depressive symptoms. The first, with follow-up of 7 to 16 years, concluded that high levels of anxiety or depression are associated with higher incidence of hypertension in white individuals aged 45 to 64 years or in black individuals aged 25 to 64 years [15]. The second, a follow up of young men for five years, also showed that high scores of depressive symptoms are associated with a higher incidence of hypertension [16].

Regarding the age group as Figure 2, most corresponded to adults, the longest phase of life, in which peak of personal achievements such as marriage, child rearing and work activities should occur. However, the presence of mental disorders in this age group often negatively influences the way of life, as it disrupts person's productivity due to their chronicity [10,17].



Some factors related to socioeconomic conditions, such as unemployment, low education, marital status, gender, poor housing conditions, informal work, and non-access to consumer goods, were identified as possible determinants to high levels of mental disorders in analyzed studies [18].

Mental health assumes an ethical-political commitment that seeks the care of people in psychological distress through respect for their uniqueness and social reintegration. Noteworthy, as part of the

therapeutic actions in the psychosocial context is drug treatment, which aims to reduce the harmful symptoms of psychiatric disorders, with the purpose of better adaptation to reality [19].

Therefore, regarding treatment, use of psychotropic drugs, such as amitriptyline and carbamazepine, was quite frequent as shown in Table 4, which agrees with the study by Sordi et al. noted that 100% of respondents who were users of a CAPS used some psychopharmaceutical. This high drug based treatment may be justified by its use in controlling and decreasing the symptoms of mental disorders, especially in those manifested by psychosis and depressed mood.

Good adherence to treatment as shown in Table 5 including psychotropic drugs may be related to significant improvement of individuals who use them. According to Xavier et al. usage of psychotropic drugs brought, in a certain way, a statement of normality for users [19]. It was observed that, sometimes, psychotropic drugs are considered an integral part of users' lives, without them they would not be able to perform their daily functions. In some moments they showed to be aware of the damage of not using, as well as revealed their expectations for cure. Therefore, therapeutic interventions should not be restricted to drug prescription.



Diagnosis	Female	Male
Epilepsy	12	13
Depression	9	1
Anxiety disorder	4	2
Schizophrenia	3	3
Psychosis	2	0
Cerebral Palsy	1	2
Panic Syndrome	3	0
Aggressive Disorder	0	1
Obsessive-compulsive disorder	0	1
Bipolarity	0	1
Alzheimer's disease	1	0
Psychomotor agitation	0	1
Behavior disorder	2	0
Autism	1	0
Parkinson's disease	1	0
Not specified	3	3
Total	42	28

Source: Research protocol

Table 4: Relationship between diagnosis and gender of patients enrolled in the USF Mental Health Program. Ananindeua-PA
*Each patient may have more than one diagnosis.

Adherence to treatment	N	%
Yes	61	98.4
No	1	1.6
Total	62	100
Source: Research protocol	3	3

Table 5: Adherence to psychoactive drug treatment by patients registered with the USF Mental Health Program. Ananindeua - PA.

Another important data was the identification of an important deficit of information in medical records. Proper and organized medical records is essential to support the direction in production of care, as it allows the planning of actions, serving as a legal instrument for the entire team of health care.

Conclusion

According to this study, the epidemiological profiles were characterized as the most prevalent syndromes, consecutively, epilepsy, depression, anxiety disorder and schizophrenia. In addition, adulthood was the most affected age group, between 30 and 39 years.

References

1. Andrade L (1999) Epidemiologia psiquiátrica. Novos desafios para o século XXI. *Revista USP* 43: 84.
2. Murray C, Lopez A (1996) The global burden of disease. (1st edtn), Harvard School of Public Health on behalf of the World Health Organization and the World Bank, Cambridge, MA, USA.
3. Penn D, Chamberlin C, Mueser K (2003) The effects of a documentary film about schizophrenia on psychiatric stigma. *Schizophrenia Bulletin* 29: 383-391.
4. Gabriela Cruz d. Crenças e atitudes estigma em relação a portadores de esquizofrenia em adolescentes estudantes da série intermediária do ensino médio: avaliação e intervenção [Mestrado]. Universidade Federal de São Paulo (UNIFESP); 2006.
5. Brasil. Lei nº 10.216, de 06 de abril de 2001. Dispõe sobre a proteção e os direitos das pessoas portadoras de transtornos mentais e redireciona o modelo assistencial em saúde mental. [Internet]. *Diário Oficial da União*. 2011 [acessado em 26 de outubro de 2019].
6. Modesto TN, Santos DN. Saúde mental na atenção básica. *Revista Baiana de Saúde Pública*.
7. Vecchia M (2006) A saúde mental no Programa de Saúde da Família: estudo sobre práticas e significações de uma equipe [Mestrado]. Universidade Estadual Paulista (UNESP).
8. Mangualde AAS, Botelho CC, Soares MR, Costa JF, Junqueira AC, et al. (2013) Epidemiological profile of patients treated in a Center for Psychosocial Care. *Mental* 10: 235-48.
9. Brasil. Relatório Final da XII Conferência Nacional de Saúde [Internet]. Conselho Nacional de Saúde. 2004 [acessado em 26 de outubro de 2019]. Disponível em: http://conselho.saude.gov.br/biblioteca/Relatorios/relatorio_12.pdf
10. Cruz DP, Sena ELS, Moreira RM, Teixeira JRB, Lira SP, et al. (2014) Clinical and epidemiological profile of patients treated in psychiatric outpatient. *Revista Cubana de Enfermería* 30: 10-20.
11. Rennó JR, Demarque R, Lobo HR, Cavalsan JP, Silva AD (2012) Saude Mental at Mulher: Transcendental Psychiatric Relocations or Cyclo Reproduction. *Revista debates em psiquiatria* 2: 6-11.
12. Teixeira PJ, Rocha FL (2007) Epidemiologic studies about association of mental disorders and metabolic syndrome. *Archives of Clinical Psychiatry* 34: 28-38.
13. Gomes FA (1996) Comorbidades clínicas em psiquiatria. (1st edtn). São Paulo: Atheneu, Brasil.
14. Sordi D, Petelincar L, Bigatto KR, Santos SG, Machado AL (2015) Comorbidities in users of a mental health service. *Portuguese Journal of Mental Health Nursing* 2: 89-94.
15. Jonas BS, Franks P, Ingram DD (1997) Are symptoms of anxiety and depression risk factors for hypertension? Longitudinal evidence from the National Health and Nutrition Examination Survey I Epidemiologic Follow-up Study. *Archives of family medicine* 6: 43-49.
16. Davidson K, Jonas BS, Dixon KE, Markovitz JH (2000) Do depression symptoms predict early hypertension incidence in young adults in the CARDIA study? *Archives of internal medicine* 160: 1495-1500.
17. Da Silva TL, Maftum MA, Kalinke LP, De Freitas Mathias TA, Ferreira AC, et al. (2015) Clinical sociodemographic profile of patients undergoing treatment in the psychiatric unit of a general hospital. *Cogitare Enfermagem* 20: 45-55.
18. Santos ÉG, Siqueira MM (2010) Prevalence of mental disorders in the adult Brazilian population: a systematic review. *J Bras Psiquiatr* 59: 238-246.
19. Da Silva Xavier M, Terra MG, Da Silva CT, De Souza Mostardeiro SC, Da Silva AA, et al. (2014) Significance of Dr. Utizano de Psychopharmacos para Individuos com Transantor Mental M Acompanhamento Ambulatory. *Escola Anna Nery Revista de Enfermagem* 18: 323-329.