

Examining the Risk Factors among Omani Women: Risk Assessment and Their Implications

Melba Sheila D'Souza*, Chandrani Isac, Ramesh Venkatesaperumal, Anandhi Amirtharaj, Shreedevi Balachandran and Anitha Thanka

Department of Adult Health and Critical Care, College of Nursing, Sultan Qaboos University, Muscat, Sultanate of Oman

Abstract

Osteoporosis is common among postmenopausal women, giving rise to morbidity and diminishing the quality of life. There is lack of information about the risk factors of osteoporosis among local Omani women. To explore the risk factors of Osteoporosis among Omani Women using a Modified Fracture Risk Assessment Tool. This is an exploratory research design using a simple random sampling. A pilot study was conducted among 35 local Omani women from the Muscat region, who completed the validated and reliable Modified Fracture Risk Assessment Tool (MFRAT). Bone Mineral Density (BMD) was measured among 11 women at moderate-high risk based on MFRAT. The data was analyzed using inferential statistics. Fourteen Omani women were at moderate-high risk for Osteoporosis fracture due to premenopausal, Vitamin D Deficiency, increased parity, increased lactation period and sedentary lifestyle. Among these five women revealed low BMD and required early intervention and support. Determinants of Osteoporosis can be elicited among premenopausal Omani women with a specific risk assessment tool. There is a need to construct a cultural sensitive risk assessment tool for Omani women for early screening and detection of Osteoporosis. Nurses can identify, provide intervention and education for Omani women at-risk of fracture.

Keywords: Fracture risk assessment; Osteoporosis; Osteopenia; Omani women; Early identification; Early screening; Bone mineral density; Nursing care; Primary prevention

Introduction

Osteoporosis (OP) is characterized by low bone mineral density (BMD) and deterioration of bone structure, resulting in an increased susceptibility to fractures of the hip, spine, and wrist [1-8]. Worldwide, approximately one-third of women aged 60-70 years and two-thirds of women aged 80 and older have osteoporosis. Lifetime risk of fracture for a 50-year-old White woman is estimated at over 70%; her risk of hip fracture alone is about 14% (National Osteoporosis Foundation [9-18]. An osteoporotic fracture occurs every three seconds, with one in three women over 50 years of age expected to be burdened with a fracture at some point [19-26]. The World Health Organization (WHO) considers osteoporosis to be second only to cardiovascular disease as a public health concern [26-29]. In the Middle East life expectancy has increased, escalating the risk factors of occurrence of Osteoporosis. Omani women have an enhanced vulnerability to the risk of fracture as a result of less exposure to Vitamin D, Calcium deficiency and sedentary life. They have an increased predisposition to develop OP, which has significant long term risks. Changes in life-style practices are likely to be influenced by these women's perception of health, choices, autonomy and control over resources. Their health behaviours are also influenced by socio-economic, cultural, religious, and gender norms. Primary care interventions are directed to identify women at increased risk before the symptoms develop, whilst primary care nurses should be vigilant to assess women with suspected fragility fractures or risk for falls. There are several risk factors which often coexist to increase risk substantially. There is an acute need for periodic fracture risk assessment, to engage in health seeking behaviours, motivation for healthy bones and prevention of osteoporosis, so as to reduce the care giver burden and health care costs. This study was done to explore the osteoporotic risk factors among Omani women for development of a cultural specific risk assessment tool.

Objective

To explore the significant risk factors of Osteoporotic fracture

among local Omani Women using a Modified Fracture Risk Assessment Tool.

Methods

This is an exploratory research design to explore the risk factors of Osteoporosis among local Omani women using a Modified Fracture Risk Assessment Tool (MFRAT) and bone mineral densitometry (BMD). We have reviewed literature and risk assessment tools for Osteoporosis or fracture among women in the Middle East and worldwide. World Health Organization (WHO) FRAX [30] tool was developed to evaluate fracture risk using 12 parameters (Black et al. [4] 2001), Osteoporosis Risk Assessment Instrument (Cadarette et al. [6] 2000) has 3 parameters, and Screening Assessment for Preventing Falls [31] has 12 parameters.

We developed a Modified Fracture Risk Assessment Tool (MFRAT) based on the standardized tools and literature search. The MFRAT has 11 parameters: Osteoporosis, Vitamin D Deficiency, Fracture, long term medications (Steroid, Anticonvulsants, Estrogen, Hormone Replacement Therapy, Heparin, Thyroxine, Vitamin D, Hypoglycemics or Calcium), family/maternal history of Osteoporosis or pathological fracture, life style, smoking cigarette or sheesha, alcohol, use of dairy products, exercise, number of children, lactating period and menstrual status. The MFRAT-English version was translated to Arabic version by linguistics and back translated. The content validity was done by 5 subject experts in the field. The reliability of the tool (r 0.81) was done among 5 local Omani women

*Corresponding author: Melba Sheila D'Souza, Assistant Professor, Department of Adult Health and Critical Care, College of Nursing, Sultan Qaboos University, Muscat, Sultanate of Oman, Tel: 00968- 98137770, E-mail: melba@squ.edu.om

Received April 18, 2013; Accepted May 20, 2013; Published May 22, 2013

Citation: D'Souza MS, Isac C, Venkatesaperumal R, Amirtharaj A, Balachandran S, et al. (2013) Examining the Risk Factors among Omani Women: Risk Assessment and Their Implications. J Nov Physiother 3: 143. doi:10.4172/2165-7025.1000143

Copyright: © 2013 D'Souza MS, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

in Muscat The tool was found to be valid, reliable, appropriate and feasible for the study. Moderate-high risk scores on the MFRAT (low: 0-9, moderate: 10-19 and high risk: 20-34) were used to select women for BMD. BMD (g/cm²) is measured by dual energy x-ray absorptiometry (DEXA) scan using a T-score of the lumbar spine/hip and/or femoral neck as the gold standard [32]. The lower the BMD score the greater the risk of fracture. There is a statistical association between poor bone density and higher probability of fracture. Osteoporosis is defined as bone mineral density that is 2.5 standard deviations or more below the mean for a young adult [32].

The inclusion criteria was Omani women aged ≥ 20 years of age (Osteoporosis has a prevalence <1% in women <45 years), voluntary participation or withdrawal from the study. The exclusion criteria are pregnant Omani women and those above 80 years. An ethical approval was obtained from the College Ethics Committee in July 2011. A pilot study using a simple random sampling was done to select thirty five (35) Omani women from the Muscat community. The selected women were explained and provided a document of the purpose, protocols and data collection method. BMD is a painless, non-invasive and involves low radiation exposure. The women were free to withdraw from the research at any time, with complete benefits of medical care, treatment or legal rights. They were aware that the BMD had minimum radiation with no known side effects compared to the benefits. An informed verbal and written consent was taken from the women who were willing to participate in the study. These women were administered the Modified Fracture Risk Assessment Tool (MFRAT). All the data were treated confidentially and stored in sealed cabinets. Bone Mineral Density (BMD) was measured among 11 women at moderate-high risk based on MFRAT. The data was analyzed using inferential statistics.

Results

Personal characteristics based on MFRAT and BMD (Table 1). Among the 35 women, only 22.9% (45.5%) were in the menopausal age (>40 years) and 34.3% (34.3%) had low body weight (<60 kg/ 125 lbs) based on the MFRAT and BMD scores respectively. 28.6% (72.7%) of the women were not regular consumers of dairy products, 14.3% (45.5%) required arm support to raise themselves from a chair and 45.7% (100%) did very little exercises or no exercises. The data shows low BMD among women with moderate-high risk factors of fracture.

Clinical characteristics based on MFRAT and BMD (Table 2). 8.6% (27.3%) of the Omani women had Osteoporosis and 20% (45.5%) had systemic illness while 17.1% (45.5%) were on long term medications based on the MFRAT and BMD scores respectively. There was 'history of fracture after 40 years' among these women (11.4%, 36.4%), their mothers (22.9%, 63.6%) and family (8.6%, 27.3%). There were significant factors governing the risk or development of Osteoporosis like predominance of pre-menopausal women (62.9%, 63.6%), Vitamin D Deficiency (37.1%, 54.5%), increased parity (22.9%, 54.5%) and longer lactating period (45.7%, 54.5%). The data shows low BMD among Omani women with moderate-high risk factors for Osteoporosis.

Association between the fracture risk factors and characteristics (Tables 1 and 2). There is a significantly high association between the fracture risk factors and long term medications (p<0.02), increased parity (p<0.02) and increased lactating period (p<0.00). There is a significant association between the fracture risk factors, Vitamin D Deficiency (p<0.07), need for arm support (p<0.07) and lack of exercises (p<0.08).

Personal Characteristics	Categories	MFRAT (35)		BMD (11)		p
		F	%	NF	%	
Age in years	20-29	13.0	37.1	1.0	9.1	0.54
	30-39	14.0	40.0	5.0	45.5	
	>40	8.0	22.9	5.0	45.5	
Weight in Kg	<60	12.0	34.3	5.0	45.5	0.58
	60-69	5.0	14.3	2.0	18.2	
	>70	18.0	51.4	4.0	36.4	
Smoking/alcohol consumption	No	35.0	100.0	11.0	100.0	-
Regular use of dairy products	Yes	25.0	71.4	3.0	27.3	0.71
	No	10.0	28.6	8.0	72.7	
Arm support to raise from a chair	Yes	5.0	14.3	5.0	45.5	0.07**
	No	30.0	85.7	6.0	54.5	
Regular exercises/mobility	Never	7.0	20.0	6.0	54.5	0.08**
	1-2 times/week	9.0	25.7	5.0	45.5	
	>3 times/week	19.0	54.3	0.0	0.0	

Table 1: Personal Characteristics among Omani women.

Clinical Characteristics	Categories	MFRAT (35)		BMD (11)		p
		No.	%	No.	%	
Osteoporosis	Yes	3.0	8.6	3.0	27.3	0.32
	No	32.0	91.4	8.0	72.7	
Vitamin D deficiency	Yes	13.0	37.1	6.0	54.5	0.07**
	No	22.0	62.9	5.0	45.5	
Systemic Illness	Yes	7.0	20.0	5.0	45.5	0.63
	No	28.0	80.0	6.0	54.5	
Long term Medications	Yes	6.0	17.1	6.0	54.5	0.02*
	No	29.0	82.9	5.0	45.5	
H/O fracture after 40 years	Yes	4.0	11.4	4.0	36.4	0.48
	No	31.0	88.6	7.0	63.6	
H/O mother's fracture after age 40	Yes	8.0	22.9	7.0	63.6	0.22
	No	27.0	77.1	4.0	36.4	
H/O Family pathological fracture	Yes	3.0	8.6	3.0	27.3	0.67
	No	32.0	91.4	8.0	72.7	
Parity/Number of Children	0	14.0	40.0	1.0	9.1	0.02*
	1 to 3	13.0	37.1	4.0	36.4	
	4 or more	8.0	22.9	6.0	54.5	
Lactating Period	<1 year	19.0	54.3	5.0	45.5	0.00*
	1-2 years	16.0	45.7	6.0	54.5	
Menstrual Status	Pre menopausal	22.0	62.9	7.0	63.6	0.35
	Menopausal	4.0	11.4	2.0	18.2	
	Menstruation	9.0	25.7	2.0	18.2	

Significant at *p<0.05, **p<0.10

Table 2: Clinical Characteristics among Omani women.

Osteoporosis classification based on MFRAT and BMD (Table 3). Among the 35 women, 14 women were in the moderate-high risk category for the occurrence of a fracture or osteoporosis according to the MFRAT. Five women showed early osteoporotic risk or marked changes on the BMD.

Discussion

Omani women with significantly low BMD results revealed they were in the premenopausal age, had longer lactation period and had increased parity. These women also had Vitamin D Deficiency and were on long term medications. Pre-menopausal women with low BMD have a significant underlying cause of risk factors [33]. There is a need for pre-menopausal osteoporotic risk assessment as one third to one half of women's bone loss can be attributed to menopause, It is important to take early measures to reduce bone loss in early middle

Diagnostic tool	Osteoporosis classification	No.	%
Fracture Risk based on BMD* N=11	Normal bone density: >-1	6.0	54.5
	Osteopenia: -1 to -2.5	3.0	27.3
	Osteoporosis: <-2.5	1.0	9.1
	Severe osteoporosis: < 2.5+fragility fracture	1.0	9.1
Fracture Risk Score based on MFRAT N=35	Low risk: 0-9	21.0	60.0
	Moderate risk: 10-19	11.0	31.4
	High risk: 20-34	3.0	8.6

*T-score is the measurement of bone mineral density (BMD) by dual energy X-ray absorptiometry scan (World Health Organization 1994)

Table 3: Osteoporosis classification among Omani women.

age women [34]. The role of Vitamin D in bone health has persistently found to be in the literature. VDD among women precipitates or exacerbates Osteopenia and Osteoporosis [35]. VDD is common in the Middle East, India, China and Japan due to the socio-economic, life-style and environmental conditions [36]. Among these determinants, the significant risk factors among Omani women are non-exposure to sunlight at an early age; complete clothing covering the body (e.g. wearing of veil to cover the head, hijab), multiparity, long period of lactation, short birth intervals, sedentary life, poor dietary intake, and exercise. Parity and lactation have detrimental effects on bone density of women in developing countries [37].

The need for arm support to raise from the chair/bed/floor, lack of exercise and poor use of dairy products was cited by majority of the Omani women with early osteoporotic changes with BMD. The relationship between decreased activity levels and bone thinning [38]. Poor exercises among the perimenopausal Omani women will increase the risk for fracture. Sedentary lifestyle has an increased tendency to sustain a fracture. Higher percentage of Jewish women with 'low or no physical activity like swimming and fitness' are at high risk for fracture compared to lower percentage among women in United States [39]. The presence of low BMD (45.5%) and moderate-high-risk scores (40%) on MFRAT show that Omani women are at risk for fracture or osteoporosis. Omani women have a predictable pattern of risk in the pre-menopausal age group, with increased parity and lactation period, Vitamin D deficiency, on long term medications, need for arm support to raise from chair, and lack of exercises. There is a need to develop a cultural specific risk assessment tool to evaluate risk of fracture among this women population for early identification, screening and intervention.

Conclusion

Increased parity, longer duration and increased frequency of lactation, premenopausal, presence of VDD and no exercises prevalent among Omani women contributed to calcium depletion and bone emptying. Nearly 45.5% of the women had low BMD showing rapid bone mass loss and risk of fracture which highlights the need for educational programs for pre-menopausal Omani women. All pre-menopausal women should have consultation and counseling about lifestyle risk factors, exercise and maintenance of adequate Calcium and Vitamin D intake [40]. Early identification of women at risk, early diagnosis and intervention can be directed at preventing the occurrence of the first fragility fracture. A cultural sensitive risk assessment tool is required for early screening, detection and referral, to motivate the local women for periodic risk assessment and engage in health seeking behaviors. Fracture risk assessment tools should incorporate information about pregnancy and duration of lactation [41].

Virtually most of the risk assessment tools reviewed are not

relevant and suitable for Arab women to identify rapid bone loss or low bone mass. Cultural sensitive specific risk assessment instruments are needed for efficient identification of women most likely to have low BMD. Nurses can further help women by providing education and support to improve the adherence to osteoporotic medication and lifestyle interventions to reduce fractures and improve quality of life among women. Majority of the Arab women remains undiagnosed and unaware of the importance of early recognition and preventive treatment of osteoporosis. Simple tools are effective in selecting women for BMD measurement and expensive specialized investigations, reducing modifiable risk factors and considering early treatment among high-risk groups. The key for prevention of osteoporosis lies in creating and using specifically designed tools for early detection and subsequently managing the underlying cause with an individualized approach.

Implications for nursing practice

Nurses and nurse educators should have an opportunity to take the lead in promoting bone health through screening and, where appropriate, intervention, referral, and patient education. Nurses must be sensitized to the need for assessing the risk for fracture or osteoporosis which will help to drastically reduce the care giver burden and ensure cost-effective measure for provision of health care. Primary care prevention is recommended with 5 universal recommendations to promote bone health [42] for women. For postmenopausal women who are not deemed high risk, the recommendation is reassessment of fracture risk every 5 years [43]. Because the prevalence of osteoporosis and fragility fracture is high and because diagnostic and therapeutic tools are available, all primary care nurses should take an active role in early identification and screening of osteoporosis, with the objective of decreasing the disease burden of osteoporosis and fracture.

Acknowledgements

We thank the College of Nursing Sultan Qaboos University and the Department of Radiology Sultan Qaboos University Hospital, subject experts who validated the tool and the women participants for their support.

References

- Allali F, Maaroufi H, Aichaoui SE, Khazani H, Saoud B, et al. (2007) Influence of parity on bone mineral density and peripheral fracture risk in Moroccan postmenopausal women. *Maturitas* 57: 392-398.
- Lim LS, Hoeksema LJ, Sherin K, ACPM Prevention Practice Committee (2009) Screening for osteoporosis in the adult U.S. population: ACPM position statement on preventive practice. *Am J Prev Med* 36: 366-375.
- Bhalla AK (2010) Management of osteoporosis in a pre-menopausal woman. *Best Pract Res Clin Rheumatol* 24: 313-327.
- Black DM, Steinbuch M, Palermo L, Dargent-Molina P, Lindsay R, et al. (2001) An assessment tool for predicting fracture risk in postmenopausal women. *Osteoporos Int* 12: 519-528.
- British Orthopaedic Association (2007) The Care of Patients with Fragility Fracture.
- Cadarette SM, Jaglal SB, Kreiger N, McIsaac WJ, Darlington GA, et al. (2000) Development and validation of the Osteoporosis Risk Assessment Instrument to facilitate selection of women for bone densitometry. *CMAJ* 162: 1289-1294.
- Cardwell CR, Abnet CC, Cantwell MM, Murray LJ (2010) Exposure to oral bisphosphonates and risk of esophageal cancer. *JAMA* 304: 657-663.
- Challen V (1997) Screening for Osteoporosis in Peri-menopausal Women: Does it Fulfill the Criteria for Medical Screening? *Radiography*, 3: 191-199.
- Clinician's Guide to Prevention and Treatment of Osteoporosis.
- Compston J, Cooper A, Cooper C, Francis R, Kanis JA, et al. (2009) Guidelines for the diagnosis and management of osteoporosis in postmenopausal women and men from the age of 50 years in the UK. *Maturitas* 62: 105-108.

11. Compston J, Rosen C (2009) *Fast Facts: Osteoporosis*. Sixth edition. Health Press Ltd, Oxford.
12. Cryer C, Patel S (2002) Primary Care Strategy for Osteoporosis and Falls and National Osteoporosis Society Falls, Fragility and Fractures.
13. Cummings SR, San Martin J, McClung MR, Siris ES, Eastell R, et al. (2009) Denosumab for prevention of fractures in postmenopausal women with osteoporosis. *N Engl J Med* 361: 756-765.
14. Zieve, Susan Storck (1994) Scientific Group on Research on the Menopause in the 1990's. WHO Technical Report Series 866, Geneva, Switzerland.
15. Dawson-Hughes B, Tosteson AN, Melton LJ 3rd, Baim S, Favus MJ, et al. (2008) Implications of absolute fracture risk assessment for osteoporosis practice guidelines in the USA. *Osteoporos Int* 19: 449-458.
16. Dornemann TM, McMurray RG, Renner JB, Anderson JJ (1997) Effects of high-intensity resistance exercise on bone mineral density and muscle strength of 40-50-year-old women. *J Sports Med Phys Fitness* 37: 246-251.
17. Shuler FD, Conjeski JM, Hamilton RL (2011) Incorporating the WHO FRAX assessment tool into nursing practice. *Am J Nurs* 111: 59-62.
18. Heinonen A, Oja P, Sievanen H, Pasanen M, Vuori I (1998) Effect of two training regimens on bone mineral density in healthy perimenopausal women: A randomized controlled trial. *J. Bone Miner. Res.* 13: 483-490
19. Holick MF (2007) Vitamin D deficiency. *N Engl J Med* 357: 266-281.
20. International Osteoporosis Foundation and International Society for Clinical Densitometry. 2011. Joint Task Force 'ISCD IOF FRAX Initiative'. Interpretation and use of FRAX in clinical practice.
21. International Osteoporosis Foundation. The facts about Osteoporosis and its impact. Lyon, France: International Osteoporosis Foundation, 2004.
22. Jang SN, Choi YH, Choi MG, Kang SH, Jeong JY, et al. (2006) [Prevalence and associated factors of osteoporosis among postmenopausal women in Chuncheon: Hallym Aging Study (HAS)]. *J Prev Med Public Health* 39: 389-396.
23. Kanis JA, Glüer CC (2000) An update on the diagnosis and assessment of osteoporosis with densitometry. Committee of Scientific Advisors, International Osteoporosis Foundation. *Osteoporos Int* 11: 192-202.
24. Kanis JA, Compston J, Cooper A, Cooper C, Francis R, et al. (2010) National Osteoporosis Guideline Group. Guideline for the diagnosis and management of osteoporosis in postmenopausal women and men from the age of 50 years in the UK. 'Invest in your Bones Campaign' of the International Osteoporosis Foundation.
25. Lim LS, Hoeksema LJ, Sherin K; ACPM Prevention Practice Committee (2009) Screening for osteoporosis in the adult U.S. population: ACPM position statement on preventive practice. *Am J Prev Med* 36: 366-375.
26. Lips P (2007) Vitamin D status and nutrition in Europe and Asia. *J Steroid Biochem Mol Biol* 103: 620-625.
27. National Guideline Clearinghouse (NGC) Guideline synthesis: Screening and risk assessment for osteoporosis in postmenopausal women. In: National Guideline Clearinghouse [Web site]. Rockville (MD): 2011
28. National Osteoporosis Foundation (NOF) America's Bone Health: the State of Osteoporosis and Low Bone Mass in Our Nation. Washington, DC: National Osteoporosis Foundation, 2002.
29. National Osteoporosis Foundation. Clinician's Guide to Prevention and Treatment of Osteoporosis. Washington, D.C.: National Osteoporosis Foundation; 2008.
30. National Osteoporosis Foundation. Osteoporosis: review of the evidence for prevention, diagnosis, and treatment and cost effectiveness analysis. *Osteoporos Int* 1998; 8 (Suppl. 4): S1-88.
31. Compston J, Cooper A, Cooper C, Francis R, Kanis JA, et al. (2009) Guidelines for the diagnosis and management of osteoporosis in postmenopausal women and men from the age of 50 years in the UK. *Maturitas* 62: 105-108.
32. Nelson HD, Haney EM, Dana T, Bougatsos C, Chou R (2010) Screening for osteoporosis: an update for the U.S. Preventive Services Task Force. *Ann Intern Med* 153: 99-111.
33. Osteoporosis-primary prevention, NICE Technology Appraisal Guideline (January 2011); Alendronate, etidronate, risedronate, raloxifene and strontium ranelate for the primary prevention of osteoporotic fragility fractures in postmenopausal women.
34. Poole KE, Compston JE (2006) Osteoporosis and its management. *BMJ* 333: 1251-1256.
35. Wall E, Walker-Bone K (2008) Use of bisphosphonates and dual-energy X-ray absorptiometry scans in the prevention and treatment of glucocorticoid-induced osteoporosis in rheumatology. *QJM* 101: 317-323.
36. [No authors listed] (2010) Management of osteoporosis in postmenopausal women: 2010 position statement of The North American Menopause Society. *Menopause* 17: 25-54.
37. U.S. Preventive Services Task Force. (2011) Screening for osteoporosis: recommendation statement. *Am Fam Physician* 83: 1197-1200.
38. University of Michigan Health System (UMHS). Osteoporosis: prevention and treatment. Ann Arbor (MI): University of Michigan Health System; 2010 Jul. 15 p.
39. Vu TT, Nguyen CK, Nguyen TL, Le BM, NguyenTrung Le D, et al. (2005) Determining the prevalence of osteoporosis and related factors using quantitative ultrasound in Vietnamese adult women. *Am J Epidemiol* 161: 824-830.
40. Werner P, Olchovsky D, Shemi G, Vered I (2003) Osteoporosis health-related behaviors in secular and orthodox Israeli Jewish women. *Maturitas* 46: 283-294.
41. Fracture Risk Assessment Tool (FRAX), World Health Organization Collaborating Centre for Metabolic Bone Diseases.
42. WHO Scientific Group on the Assessment of Osteoporosis at Primary Health Care Level, World Health Organization (2004).
43. (1994) Assessment of fracture risk and its application to screening for postmenopausal osteoporosis. Report of a WHO Study Group. *World Health Organ Tech Rep Ser* 843: 1-129.

Citation: D'Souza MS, Isac C, Venkatesaperumal R, Amirtharaj A, Balachandran S, et al. (2013) Examining the Risk Factors among Omani Women: Risk Assessment and Their Implications. *J Nov Physiother* 3: 143. doi:[10.4172/2165-7025.1000143](https://doi.org/10.4172/2165-7025.1000143)

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 250 Open Access Journals
- 20,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission/>