

Euro Surgery 2020: Management of Acne Scarring

Name: Chang D

Medical Director, Academy of Aesthetic Medicine, Korean Medical Aesthetics Clinic, Singapore

Short Communication: Acne scarring is a fibrous process involving new collagen production to heal a wound. It affects those with acne vulgaris and is predominant in nodulocystic acne [1]. Options for treatment depends on acne scarring severity, scar type (Table 1), scar site and patient skin type, choice of treatment modality and side effects risk profile. This paper attempts to share with you the management of acne scarring [2].

Different types of scars	Symptoms
Flat scars	These can be red, brown or pale flat marks
Mild hypertrophic and atrophic scars	Sunken or raised scars, not visible at distances of 50cm or more
Moderate hypertrophic and atrophic scars	Visible at distances of ≥ 50 cm, camouflaged and can be even out by stretching the skin
Severe hypertrophic and atrophic scars	Visible at distances ≥ 50 cm, difficult to be camouflaged and cannot be even out by stretching of the skin

Unfortunately, hypertrophic or keloid scars tend to reappear even after a good result is obtained post treatment. Traditionally, scar revision surgery is employed. Ablative lasers, carbon dioxide and erbium YAG laser are successful for surgery. To improve the color of scars, Pulse Dye Laser (PDL) has been effectively utilized. The PDL can reduce erythema and textures of the scar. Multiple treatments are needed.

A microneedle radiofrequency laser, an example of which is the INTRAcel from Korea, has produced significant results. It uses an insulated microneedle that delivers radiofrequency energy to the dermis,

reducing damage to the epidermis, hence speeding up recovery. In my practice, I use a schedule of 4 sessions of INTRAcel, and can achieve results of 60% upwards for acne scars. More recently, the plasma laser have been used, achieving good results with minimal downtime and epidermal damage.

Ultimately the challenge is true acne scars never completely fade, although they visibly improve as time passes. The good thing is, they can be concealed. Patient skin type is also an important factor. Skin type can be classified from Fitzpatrick 1 to 6. Darker skin types have a higher risk of post inflammatory hyperpigmentation. Hence they benefit more from fractional laser compared to traditional full ablative lasers. Scars can occur anywhere on the body, although the upper chest and shoulders are especially prone to them. This is due to poorer regeneration capability, from the lower density of adnexa structures (sebaceous glands, hair follicles).

Technology and Treatment Factors

The major complication of acne is scarring, which results in severe comorbidity for patients. On the bright side, there exists several options to treat acne scarring. Studies support combination treatment to achieve the best results. The safety and efficacy has good evidence, but there is a dearth of split-face trials [3].

Scar types and treatment

- Grade 1 scars: Topical treatment with silicone based gels combined with careful use of pigment, vascular and fractional lasers.
- Grade 2 mild scars: Fractional, vascular and pigment lasers as well as microneedling.

- Grade 3 scars: Full ablative and fractional lasers, and may require scar revision occasionally.
- Grade 4 scars: Require a multimodality approach [4].

Ablative laser: Ablative laser resurfacing utilizes electromagnetic waves to erode the entire epidermis and upper layer of dermis. Fractional laser treatment addresses both the epidermis and dermis. It delivers hundreds of laser beamlets. The principle is controlled dermal damage in MTZ (Micro Thermal Treatment Zones) in order to stimulate fibroblast to aid collagen remodeling and neocollagenesis. MTZ are selected areas with dermal injury, while sparing adjacent areas. They areas with injury would in turn kick start the process of neocollagenesis, while the adjacent areas maintain cell renewal and healing. In this way, the electromagnetic energy is effective yet safe, meanwhile keeping surrounding tissue healthy, aiding wound healing. This leads to faster recovery and less downtime. Today, the main laser resurfacing treatments can be divided into the non-ablative and the fractional. Techniques using other modalities may include microneedling, high intensity focused ultrasound systems and platelet rich plasma amongst others [5].

Non ablative laser

Non ablative lasers deliver electromagnetic energy into the dermis, inducing selective injury, leaving the epidermis untouched. The safety profile and speedy recovery have more than compensated for its slower results. And in today's day and age, the public want treatments that have minimal downtime. There are 3 main types of non-ablative modalities, mid-infrared ones, the Pulse dye lasers (PDL) and broadband light systems. These lasers work well for dyschromia but the textural improvement in scars is minimal [6].

CO2 ablative laser was the primary modality for the treatment of acne scars. However, it has extensive downtime of 2 or more weeks. Erbium YAG laser has slower clinical efficacy, shorter downtime and minimal recovery time. Now, the focus is on multi-modality treatments to achieve good results, safe and effective with low downtime. Hence the popularity of fractional lasers [7-9].

Side Effects: Side effects include redness and swelling, skin dyschromia, scarring, skin infections, eczema etc. After ablative fractional lasers, these following are common side effects, especially over the neck.

To bring it to another level, RF microneedling is a new technique invented in Korea. It works to reduce scars, both atrophic and hypertrophic scars. The science is dermal matrix remodeling. The downtime is minimal; must less than fractional lasers, the side effects of post inflammatory increased pigmentation on darker skin types is much reduced, and the results are comparable [16].

Dermal fillers: A soft filler like restylane vital can be used to fill up the depression in atrophic scars, and subcision can be employed to breakdown the fibers that are anchoring the skin down. In my practice, I commonly combine this with RF Microneedling or fractional laser, over 4 sessions, and the results are significant. Studies have noted the use of PRP can speed up recovery [17].

Plasma laser: This relies on the 4th state of matter to ionize particles. It removes oxygen from cells, hence reducing the risk of charring the skin. The heat reaches a good temperature to stimulate neocollagenesis to improve scar texture and tone. The safety profile is high, with low risk of hypopigmentation and hyperpigmentation.