



Neutrophil / Lymphocyte Ratio (NLR) – Trombocyte / Lymphocyte Ratio (TLR): A Predictor of Axillary Lymph Node Metastasis in Breast Cancer Patients?

Tugba Han Öner

Baskent University School of Medicine, Turkey

Abstract:

Background: This study evaluated the relationship between preoperative neutrophil / lymphocyte ratio - platelet / lymphocyte ratio, clinicopathological, radiological factors, and axillary lymph node metastasis in stage I-III breast cancer to determine if axillary surgery can be safely omitted in selected patients.

Methods: The study included 158 Stage I-III breast cancer patients operated on at Baskent University Zubeyde Hanım Research Center between 2011 and 2018. The incidence of axillary lymph node metastasis was correlated with clinical, radiological, pathological, and laboratory (neutrophil count to lymphocyte count, platelet count to lymphocyte count) findings by univariate and multivariate analyses. Sensitivity and specificity calculations, positive predictive value, negative predictive value, positive and negative Likelihood Ratio (accuracy ratio), and exact accuracy were calculated for neutrophil/lymphocyte ratio cut-off values of 3.5 and 1.

Results: Neutrophil and platelet values were significantly higher in patients with lymph node metastasis. Neutrophil / lymphocyte ratio - platelet / lymphocyte ratio values were higher in patients with axillary lymph node metastasis, but this was not statistically significant. Axillary lymph node metastasis was not associated with age, lymphocyte, monocyte count, estrogen receptor, progesterone receptor, or c-erb B2 status. The incidence of axillary lymph node metastasis was statistically significantly higher in the presence of lymphovascular invasion. Sensitivity, specificity, positive predictive value, and negative predictive value were 93.85%, 16.67%, 44.9%, and 78.9% respectively for axillary lymph node metastasis while the neutrophil / lymphocyte ratio was ≥ 3.5 . Specificity, sensitivity, positive predictive value, and negative predictive value were 97.78%, 9.23%, 75.0%, and 59.9%



respectively for axillary lymph node metastasis while the neutrophil / lymphocyte ratio < 1 .

Conclusions: For axillary lymph node metastasis, neutrophil, platelet counts, lymphovascular invasion status, radiological and pathological mass size, and presence of radiological axillary lymphadenopathy are the statistically significant independent variables. They provide information that can help surgeons decide on the treatment of breast cancer patients with certain neutrophil / lymphocyte ratio values (neutrophil / lymphocyte ratio < 1 and neutrophil / lymphocyte ratio ≥ 3.5).

Biography:

Tugba Han Öner works as a General Surgery specialist in İzmir Baskent University Zübeyde Hanım Application and Research Center in İzmir Karşıyaka district.

Publication of speakers:

1. Ballıkaya, Sedat & Oner, Yildirhan & Temel, Tugba & Ozkal, Burak & Bailey, Trevor & Toprak, Muhammet & Uher, Ctirad. (2019). Thermoelectric and thermal stability improvements in Nano-Cu₂Se included Ag₂Se. *Journal of Solid State Chemistry*. 273. 10.1016/j.jssc.2019.02.037.
2. Oner, Ayse & Capraro, Robert & Capraro, Mary. (2016). The Effect Of T-STEM Designation On Charter Schools: A Longitudinal Examination Of Students' Mathematics Achievement. *Sakarya University Journal of Education*. 6. 80. 10.19126/suje.17778.

[International Conference on Surgery and Anesthesia | August 10, 2020 | London, UK](#)

Citation: Tugba Han Öner; Neutrophil / Lymphocyte Ratio (NLR) – Trombocyte / Lymphocyte Ratio (TLR): A Predictor of Axillary Lymph Node Metastasis in Breast Cancer Patients?; Euro Surgery 2020: August 10, 2020; London, UK