

Editorial

Open Access

Peptide Coupling Reactions

Khattab NS¹, El-Faham A^{1,2*}, and Albericio F²⁻⁷

¹Chemistry Department, Faculty of Science, Alexandria University, P.O. Box 426, Ibrahimia, 12321 Alexandria, Egypt

²Department of Chemistry, College of Science, King Saud University, P.O. Box 2455, Riyadh 11451, Saudi Arabia

³Institute for Research in Biomedicine (IRB), Barcelona Science Park, Baldíri Reixac 10, Barcelona 08028, Spain

⁴CIBER-BBN, Networking Centre for Address, Baldíri Reixac 10, 08028-Barcelona, Spain

⁵Department of Organic Chemistry, University of Barcelona, Martí i Franqués 1-11, Barcelona 08028, Spain

⁶School of Chemistry & Physics, University of KwaZulu-Natal, Durban 4001, South Africa

⁷School of Chemistry, Yachay Tech, Yachay City of Knowledge, 100119-Urcuqui, Ecuador

Peptide bond formation is a nucleophilic substitution reaction of an amino group (nucleophile) at a carboxyl group involving a tetrahedral intermediate. Furthermore, the peptide coupling reaction must be performed under mild conditions, and preferably at room temperature. Activation of the carboxyl component is achieved by the introduction of electron accepting moieties [1]. Carboxyl components can be activated as acyl halides, acyl azides, acylimidazoles, anhydrides, esters etc. There are different ways of coupling reactive carboxyl derivatives with an amine [2].

In recent years, peptide-coupling reactions have significantly advanced in accord with the development of new peptide-coupling reagents and their application to both solution and solid-phase synthesis [3,4]. The formerly techniques of carbodiimide is being replaced with iminium/uronium derivatives 1-5 [5-21] (Figure 1).

Later, El-Faham and Albericio [22] reported a new family of coupling reagents based on the modification of the structures of the carbocation skeleton moiety, which feature relatively high reactivity and low racemization during peptide bond formation [22]. Very recently, El-Faham and Albericio [23] extended their work taking an N-containing 6-membered ring structure containing O, S, and N-CH₃ for synthesis of novel coupling reagents [24] (Figure 2).

Recent reports confirmed the explosive properties of HOBT derivatives [25]. Accordingly, El-Faham and Albericio [25-26] reported the new additives as well as their uronium salts derivatives as replacement for HOBT and HOAt derivatives (Figures 3 and 4). Among these entire additives Oxyma [26] (Figure 3) and its uronium salt COMU (Figure 4) showed an excellent replacement for HOBT and its analogues [26,27].

More recently, we have reported 5-(hydroxyimino)-1,3-dimethylpyrimidine-2,4,6 (1H,3H,5H)-trione (Oxyma-B) as an excellent additive for the suppression of racemization during peptide




Figure 2: Structure of morpholine-based coupling reagents.




Figure 3: Structure of oximes to replace of HOBT.

synthesis [27]. Oxyma-B, has the same structure future for the carbonyl moiety in which the oxime group is flanked between the two carbonyl group as in HONM. In addition, Oxyma-B performs better as a racemization suppressor than Oxyma Pure and even better than HOAt in both stepwise and segment coupling in solid- and solution-phase peptide synthesis [28]. Lately, a new class of O-form uronium-type coupling reagents derived from Oxyma-B were introduced TOMBU and COMBU (Figure 5) [29].






Figure 1: Structure of symmetric iminium/uronium salts coupling reagents.

*Corresponding author: El-Faham A, Chemistry Department, Faculty of Science, Alexandria University, P.O. Box 426, Ibrahimia, 12321 Alexandria, Egypt, E-mail: aymanel_faham@hotmail.com

Received February 07, 2015; Accepted February 09, 2015; Published February 15, 2015

Citation: Khattab NS, El-Faham A, Albericio F (2015) Peptide Coupling Reactions. J Mol Pharm Org Process Res 3: e119. doi: 10.4172/2329-9053.1000e119

Copyright: © 2015 Khattab NS, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited



References

- Montalbetti CAGN, Falque V (2005) Amide bond formation and peptide coupling. *Tetrahedron* 61: 10827-10852.
- El-Faham A, Albericio F (2011) Peptide Coupling Reagents, More than a Letter Soup. *Chem Rev* 111: 6557-6602.
- Albericio F, Carpino LA (1997) Methods Enzymol: Solid Phase Peptide Synthesis Fields, Academic Press, Orlando 289: 104-126.
- Albericio F, Kates SA (2000) In Solid-Phase Synthesis, A Practical Guide, Marcel Dekker, New York 275-330
- Albericio F, Chinchilla R, Dodsworth DJ, Najera C (2001) New Trends in Peptide Coupling Reagents. *Org Prep Proced Int* 33: 203-303.
- Humphrey JM, Chamberlin AR (1997) Chemical Synthesis of Natural Product Peptides: Coupling Methods for the Incorporation of Noncoded Amino Acids into Peptides. *Chem Rev* 97: 2243-2266.
- Valeur E, Bradley M (2009) Amide bond formation: beyond the myth of coupling reagents. *Chem Soc Rev* 38: 606-631.
- König W, Geiger RA (1970) New method for the synthesis of peptides: activation of the carboxyl group with dicyclohexylcarbodiimide and 1-hydroxybenzotriazoles. *Chem Ber* 103: 788-798.
- Carpino LA (1993) 1-Hydroxy-7-azabenzotriazole. An efficient peptide coupling additive. *J Am Chem Soc* 115: 4397-4398.
- Knorr R, Trzeciak A, Bannwarth W, Gillessen D (1989) New coupling reagents in peptide chemistry. *Tetrahedron Lett* 30: 1927-1930.
- Abdelmotti I, Albericio F, Carpino LA, Forman BM, Kates SA (1994) Structural studies of reagents for peptide bond formation: Crystal and molecular structures of HBTU and HATU. *Lett Pept Sci* 1: 57-67.
- Dourtoglou V, Ziegler JC, Gross B (1978) O-Benzotriazolyl-N,N-tetramethyluronium hexafluorophosphate: a new and effective reagent for peptide coupling. *Tetrahedron Lett*: 1269-1272.
- Wijkmans JCHM, Blok FAA, Van der Marel GA, Van Boom JH, Bloemhoff W (1995) CF₃-NO₂-PyBOP: a new and highly efficient coupling reagent for N-methyl amino acids. *Tetrahedron Lett* 36: 4643-4646.
- Carpino LA, El-Faham A, Albericio F (1995) Efficiency in Peptide Coupling: 1-Hydroxy-7-azabenzotriazole vs 3, 4-Dihydro-3-hydroxy-4-oxo-1,2,3-benzotriazine. *J Org Chem* 60: 3561-3564.
- Klose J, El-Faham A, Henklein P, Carpino LA, Bienert, M (1999) Addition of HOAt Dramatically Improve the Effectiveness of Pentafluorophenyl-Based Coupling Reagents. *Tetrahedron Lett* 40: 2045-2048.
- Chen SQ, Xu JC (1992) A new coupling reagent for peptide synthesis. Benzotriazolylloxy-bis (pyrrolidine)-carbonium hexafluorophosphate (BBC). *Tetrahedron Lett* 33: 647-650.
- Carpino LA, El-Faham A (1994) Effect of Tertiary Bases on O-Benzotriazolyluronium Salt-Induced Peptide Segment Coupling. *J Org Chem* 59: 695-698.
- Carpino LA, Xia J, El-Faham A (2004) 3-Hydroxy-4-oxo-3, 4-dihydro-5-azabeno-1,2,3-triazene. *J Org Chem* 69: 54-61.
- Carpino LA, Henklein P, Foxman BM, Abdelmotti I, Costisella B et al. (2001) The Solid and Solution Structure of HAPyU. *J Org Chem* 66: 5245-5247.
- Li P, Xu JC (2000) New and Highly Efficient Immonium Type Peptide Coupling Reagents: Synthesis, Mechanism and Application. *Tetrahedron* 56: 4437-4445
- Li P, Xu JC (2000) HOBT and HOAt-derived immonium salts: new and highly efficient coupling reagents for peptide synthesis. *Tetrahedron Lett* 41: 721-724.
- El-Faham A, Khattab ShN, Abdul-Ghani M, Albericio F (2006) Design and Synthesis of New Immonium-Type Coupling Reagents. *Eur J Org Chem* 1563-1573.
- El-Faham A, Albericio F (2007) Novel Proton Acceptor Immonium-Type Coupling Reagents: Application in Solution and Solid Phase Peptide Synthesis. *Org Lett* 9: 4475-4477.
- Faham A, Albericio F (2008) Morpholine-Based Immonium and Halogenoamidinium Salts as Coupling Reagents in Peptide Synthesis. *J Org Chem* 73: 2731-2737.
- Wehrstedt KD, Wandrey PA, Heitkamp D (2005) Explosive properties of 1-hydroxybenzotriazoles. *J Hazard Mater* 126: 1-3.
- Subiros-Funosas R, Prohens R, Barbas R, El-Faham A, Albericio F (2009) Oxyma: an efficient additive for peptide synthesis to replace benzotriazole-based HOBT and HOAt with a lower risk of explosion. *Chem Eur J* 15: 9394-9403.
- El-Faham A, Subirós-Funosas R, Prohens R, Albericio F (2009) COMU, a Safer and More Effective Replacement for Benzotriazole-based Uronium Coupling Reagents. *Chem Eur J* 15: 9404-9416.
- Jad YE, Khattab ShN, Govender Th, Kruger HG, El-Faham A, et al. (2014) Oxyma-B, an Excellent Racemization Suppressor. *Org and Bio Chem* 12: 8379-8385.
- Jad YE, Khattab ShN, de la Torre BG, Govender Th, Kruger HG, et al. (2014) TOMBU and COMU as novel uronium-type peptide coupling reagents derived from Oxyma-B. *Molecules* 19: 18953-18965.