



Research Article

SYNTHESIS, CHARACTERIZATION AND BIOLOGICAL STUDIES OF Fe(II), Ce(II) AND Cr(II) COMPLEXES WITH N,N'- BIS(BENZOIN)-1,4 BUTANE DIIMINE

V. Balamurugan^{1*}, S. Sankar²

1. Department of Chemistry, Saranathan College of Engineering , Tiruchirapalli, Tamil Nadu, India
2. Department of Chemistry, A.V.V.M. Sri Pushpam College, Thanjavur, Tamil Nadu, India

*Corresponding Author: Email manibala_mb@yahoo.co.in

(Received: August 17, 2014; Accepted: October 30, 2014)

ABSTRACT

Objective: The objective of the present study is to synthesize Schiff base ligand, N,N' -bis(benzoin) 1,4 butane diimine and its Fe(II), Co(II) and Cr(II) complexes.

Methods: The synthesized complexes were characterized by IR and SEM. The synthesized Schiff base ligand and its Fe(II), Co(II) and Cr(II) complexes were screened for antibacterial and antifungal activity.

Results: IR data show that the complexes are four coordinate. The SEM show that all the complexes are nano crystalline.

Conclusion: The synthesized complexes were tested against the bacterial and fungal species. The synthesized complexes were showed remarkable zone of inhibition with standards and have good antibacterial and antifungal activities.

Keywords: Schiff base, N,N' -bis (benzoin) 1, 4 butane diimine, Metal Complex, Antimicrobial.

INTRODUCTION

The reaction of primary amines and an aldehyde or a ketone forms a Schiff base. A Schiff base is a chemical compound containing of a -C=N- group [1-3]. Transition metal Schiff base complexes have been generally prepared by refluxing the solutions of a metal salt, a diamine and a compound containing a carbonyl group and a primary amine. Holm et al. (1966) reported the synthesis and characterization of some divalent copper complex compounds by refluxing the solutions of copper (II) salt, salicylaldehyde but different diamines. Thakur et al. [4] reported the synthesis and characterization of Ni (II), Co (II) and Fe (II) Schiff base complex compounds. The compounds were prepared by refluxing mixture of the solutions of a divalent metal chloride, acetylacetone and hydrazine hydrate.

However, recent reports indicated synthetic modification in which the ligand is separately prepared before its interaction with a metal salt. Xishi et al., 2003[5] showed the synthesis and spectroscopic properties of Mn (II), Co (II) and Cu (II) complexes with novel Schiff base ligand derived from 2, 2' bis(p-methoxyphenylamine) and salicylic aldehyde. Schiff base complexes of transition metals are of great importance in medicine, biochemistry and industries among others. For example, the field of medicine has witnessed an increase in the number of complexes with therapeutic value. Cobalt (II) Schiff base complexes are potential antiviral agents, cis-dichlorodiamineplatinum (II) complex is an anti cancer agent while copper (II) Schiff base is an anti-tubercular agent [6, 7]. This paper reports the synthesis and characterization of N, N' – Bis (Benzoin)- 1,4 butane diimine metal(II) complex compounds due to paucity of information.

MATERIAL AND METHOD

The chemicals and solvents used in this work were of Analar grade. All the glass wares used were washed thoroughly with distilled water and dried in an oven. Infrared spectra were recorded on a Perkin-Elmer FT-IR type 1650 spectrophotometer in wave number region 200-4000 cm^{-1} . SEM images were recorded in VEGA3 TESCAN SEM analyzer.

Preparation of N, N' – Bis(Benzoin)- 1,4 butane diimine

To a methanolic solution of 2.12g (0.01mol) benzoin and 1.09g (0.005mol) 1,4 butane diimine was added anhydrous sodium acetate (4g) and the mixture refluxed for an hour. The hot solution was poured into ice-cold water where upon yellow precipitate of the schiff's base separated, it was filtered. Washed with water, dried and recrystallised from ethanol[8].

Preparation of N, N' – Bis(benzoin)- 1,4 butane diimine Metal (II) Complexs

The ethanolic mixture of each(0.01mol) Iron (II) chloride, (0.01mol) Cobalt (II) chloride, Chromium(II) chloride, and N, N' – bis(benzoin) butanediiimine schiff base (0.01mol) was refluxed separately, followed by drop wise addition of ammonia. The metal chelate separated out was filtered, washed with ethanol followed by ether and dried in an oven at 50°C [8].

Determination of antimicrobial activity

The antimicrobial activity was performed by agar cup plate method.

Microorganisms

Escherichia coli (Gram negative) and fungi as Candida albicans were the microorganisms used and they were obtained from the Microbiology Laboratory of the Thanjavur Medical College Hospital, Thanjavur. These microorganisms were identified and confirmed by Microbiologists, Department of Microbiology, Thanjavur Medical College, Thanjavur.

Preparation of 24 hours pure culture

A loop full of each of the microorganisms was suspended in about 10ml of physiological saline in a Roux bottle. Each of these was streaked on to the appropriate culture slants and was incubated at 37°C for 24 hours except for Candida albicans which was incubated at 25°C for 24-48 hours. After

completion of incubation period, when growth was observed the tubes were kept into 2-8°C until use.

Sample solutions for the experiment

The sample solutions as 50 μl , 100 μl and 150 μl were used for the experiment. Standard antibiotic solution as Chloromphenical for bacteria and Nystatin (25mg/ml distilled water- 30 μl) for fungi used to compare the test solution.

Preparation of dried filter paper discs

Whatman filter paper (No:1) was used to prepare discs approximately 6 mm in diameter, which are placed in hot air for sterilization. After sterilization, the discs were loaded with different concentrations of prepared sample solutions again kept under refrigeration for 24 hrs.

Application of discs to inoculated agar plates

Previously prepared paper discs were dispensed onto the surface of the inoculated agar plate. Each disc was pressed down firmly to ensure complete contact with the agar surface. The discs were placed on the medium suitably apart and the plates were incubated at 5°C for 1 hr to permit good diffusion and then transferred to incubator at 37°C for 24 hrs. After completion of 24hrs, the plates were inverted and placed in an incubator set to respective temperature for 24 hrs

Antimicrobial assay

Antibiogram was done by disc diffusion method [9,10]. Petri plates were prepared by pouring 30 ml of NA /PDA medium for bacteria/fungi. The test organism was inoculated on solidified agar plate with the help of micropipette and spread and allowed to dry for 10 mins. The surfaces of media were inoculated with bacteria/fungi from a broth culture. A sterile cotton swab is dipped into a standardized bacterial/ fungi test suspension and used to evenly inoculate the entire surface of the Nutrient agar/PDA plate. Briefly, inoculums containing Escherichia coli was spread on Nutrient agar plates for bacteria and Candida albicans was spread on potato dextrose agar for fungus strains. Using sterile forceps, the sterile filter papers (6 mm diameter) containing the sample (50 μl , 100 μl , 150 μl and 30 μl for respective standard) were laid down on the surface of inoculated agar plate. The plates were incubated at 37°C for 24 h for the bacteria and at room temperature (30 \pm 1) for 24-48 hr. for

yeasts strains. Each sample was tested in triplicate.

Measurement of zone of inhibition

The antimicrobial potential of test compounds was determined on the basis of mean diameter of zone of inhibition around the disc in millimeters. The zones of inhibition of the tested microorganisms by the extracts were measured using a millimeter scale. The diameter sizes in mm of the zone of inhibition are shown in the table 2 to 4 and Fig. 2 to 4.

RESULTS AND DISCUSSION

The interaction between metal (II) ions and N, N' – Bis(Benzoin)- 1,4 butane diimine, yielded the desired crystalline, yellow metal (II) complexes compound. These compounds are insoluble in water and common organic solvents, but are readily soluble in acetone.

The IR spectra [Table 1] of the metal (II) complex compounds show bands in the region 1530 - 1570cm⁻¹ which are assigned to $\nu(\text{C}=\text{N})$ stretching vibration, a fundamental

Table 1: The Infrared Spectral Data of the Complex Compounds.

Compound	$\nu(\text{C-H}) (\text{cm}^{-1})$	$\nu(\text{C=C}) (\text{cm}^{-1})$	$\nu(\text{C=N}) (\text{cm}^{-1})$	$\nu(\text{M-O}) (\text{cm}^{-1})$	$\nu(\text{M-N}) (\text{cm}^{-1})$
[Fe L']	2822	1610	1569	580	465
[Co L']	2942	1612	1567	586	458
[Cr L']	2892	1614	1560	581	460

Table 2: Antimicrobial activity of Ligand

Ligand	<i>Escherchia Coli</i> (mm)	<i>Candida albicans</i> (mm)
50 μ l	-	2 \pm 0.14
100 μ l	8 \pm 0.56	4 \pm 0.28
150 μ l	13 \pm 0.91	5 \pm 0.35
Standard (30 μ l) (Chloromphenical)	13 \pm 0.91	-----
Standard (30 μ l) (Nystatin)	-----	13 \pm 0.91

Table 3: Antimicrobial activity of Iron complex

Cobalt complex	<i>Escherchia Coli</i> (mm)	<i>Candida albicans</i> (mm)
50 μ l	-----	-----
100 μ l	-----	-----
150 μ l	12 \pm 0.84	10 \pm 0.7
Standard (30 μ l) (Chloromphenical)	12 \pm 0.84	-----
Standard (30 μ l) (Nystatin)	-----	14 \pm 0.98

Table 4 (a): Antimicrobial activity of Cobalt complex

Cobalt complex	<i>Escherchia Coli</i> (mm)	<i>Candida albicans</i> (mm)
50 μ l	-----	-----
100 μ l	-----	-----
150 μ l	12 \pm 0.84	10 \pm 0.7
Standard (30 μ l) (Chloromphenical)	12 \pm 0.84	-----
Standard (30 μ l) (Nystatin)	-----	14 \pm 0.98

Table 4 (b): Antimicrobial activity of Chromium complex

Chromium complex	<i>Escherchia Coli</i> (mm)	<i>Candida albicans</i> (mm)
50 μ l	-----	-----
100 μ l	-----	-----
150 μ l	10 \pm 0.84	-----
Standard (30 μ l) (Chloromphenical)	12 \pm 0.84	-----
Standard (30 μ l) (Nystatin)	-----	14 \pm 0.98




Fig 2: SEM images of Fe(II), Co(II) and Cr(II) Complexes

feature of azomethine group[11].

The band in the region 1600 - 1660cm⁻¹ corresponds to $\nu(\text{C}=\text{C})$ stretching vibration due to phenyl group, indicating the coordination of bis(benzoin)-1,4 butane diimine Schiff base to the metal (II) ions. The bands within 550 - 590cm⁻¹ and 430 - 480cm⁻¹ are assigned to $\nu(\text{M}-\text{O})$ and $\nu(\text{M}-\text{N})$ stretching vibrations, respectively [12,13]. These bands confirmed the coordination of the Schiff base to the metal (II) ions. The metal (II) complex compounds revealed 1:1 metal to ligand ratio, resulting in a four coordinate complex compounds of which the following molecular structure below is proposed.




Fig1: Shows the molecular structure of bis (benzoin)- 1, 4 butane diimine complex compounds

Scanning Electron Microscope

The SEM micrographs of Zn(II) and Nickel(II) complexes are shown in Figs.2a, 2b and 2c. It is seen from the figure that Zn(II) and Nickel(II) complexes shows cauliflower like structures. The particle sizes of Zn (II) and Nickel (II) complexes were in the diameter range of few microns. However, particles with size less than 100nm were also observed which groups agglomerates of larger size. The average crystalline size shows that the particles were agglomerated that these complexes were polycrystalline with nanosized grains.

CONCLUSION

The Schiff base ligand and its Fe(II), Co(II) and Cr(II) complexes were prepared. Fe(II), Co(II) and Cr(II) complexes were characterized by IR spectra and SEM. The proposed structure of metal complexes has been illustrated in Fig.1. The surface morphology using SEM showed that the particles were polycrystalline with nano sized grains. The antimicrobial activity of ligand and its complexes indicate that both ligand and metal complexes are potentially active against both fungi and bacteria.

Acknowledgments

The authors are thankful to Mr. S.Chandra Mohan, Asst.Professor, Department of Chemistry, PRIST University, Thanjavur, Dr.S.Velavan, Harman Research Centre, Thanjavur and Mr. R.Ramadoss, Periyar TBI, Periyar Maniammai University, Vallam, Thanjavur for providing research facilities.

REFERENCES:

- Holm RH, Everett JR, Chakravorty RS. Metal Complexes of Schiff Bases and β -ketoimines. *Inorganic Chemistry* 1996; 7: 83-214.
- Gupta Hobday MD, Smith TD. N, N'-ethylene bis(salicylideneiminato) Transition Metal ion Chelates. *Coordination Chemistry Review* 1972; 9: 311-337.
- Pierre. *Organic reactions*, Johnson Wiley Pub. New York USA 1987. p. 73-79.
- Thakur RS, Sharman RU, Rai HC. Studies on Ni (II), Co (II) and Fe (II) Schiff base complexes, *Indian Journal of Chemistry* 1982; 21A: 837-838
- Xishi T, Xian HY, Qiang, C, Minyu T. Synthesis of Some Transition Metal Complexes of a Novel Schiff Base Ligand Derived from 2, 2' - Bis(p-Methoxy Phenylamine) and Salicylic Aldehyde. *Molecules* 2003, 8: 439-443.
- Lippard SJ. *Bioinorganic Chemistry*, 2nd Edition, University Science Books Millvalley; 1996. p. 505-508.
- Bleomink MJ, Reedi JK. *Metal Ions in Biological Systems*, 3rd Edition, MerceL Dekker Inc. New York; 1996. p. 641-644.
- Mohapatra BB, Mohapatra, Guru S. J. *Inorganic, Nuclear Chemistry* 1978, 40 : 1179-1180.
- NCCLS-National Committee for Clinical Laboratory Standards. *Performance standards for antimicrobial disc susceptibility tests*. PA: NCCLS Publications; 1993, p. M2-A5.
- Chittibabu.B., Ayya R.M., "Determination of major organic constituents in FK 506 treated tissues of albino rats", *Int. J. Res. Dev.Pharm. L. Sci.*, 2014, 3(5), pp. 1151-1162.
- Awoyinka O, Balogun IO, Ogunnowo AA. Phytochemical screening and in vitro bioactivity Of *Cnidocolus aconitifolius* (Euphorbiaceae). *J Med Plant Res* 2007; 1(3): 63-65.
- Boucher LJ, Day WV. *Electrochemical Method*, 2nd Edition, Willey New York; 1977. p. 6-13
- Nakomato K. *Infrared Spectra of Inorganic and Coordination Compounds*, Willey, New York; 1963 .p. 234-241.

How to cite your article:

Balamurugan V., Sankar S., "Synthesis, characterization and biological studies of fe(ii), ce(ii) and cr(ii) complexes with N,N'-bis(benzoin)-1,4 butane diimine", *Int. J. Res. Dev. Pharm. L. Sci.*, 2015, 4(1), pp. 1323-1327.