

The Ascidian *Ciona Intestinalis* as Model Organism for Ecotoxicological Bioassays

Alessandra Gallo and Elisabetta Tosti*

Department of Biology and Evolution of Marine Organisms, Stazione Zoologica Anton Dohrn, Napoli, Italy

Editorial

The increase of pollutants in the oceans as well as the global change around the world is having a negative impact on the biodiversity of living organisms. There is a crescent alarm in the scientific community for the different types of pollution (biological, chemical and physical) and for the changes occurring at the level of species and communities. Many parameters endangering marine wildlife contribute to create an unhealthy marine environment that induces reproductive disorders that, in turn, may influence even the survival of marine wildlife.

Ascidians, or sea squirts, are sessile marine invertebrates commonly found throughout the world. They have been largely used as models for developmental biology since the last century [1]. The advantage of using ascidians for studies of reproduction and development is their large number of oocytes, their synchronous arrest in the metaphase of the first meiotic division, the rapid embryo development that generates swimming larvae around 18-24 hr post-fertilization, and the small number of cells and few tissues in the adult. *Ciona intestinalis* is among the most studied ascidian, and its genome is well established since from 2003 [2].

From an experimental point of view, *Ciona* oocytes are relatively easy to manipulate and microinject, this further advantage has rendered this species a model to study reproductive mechanisms and meiotic regulation [3-6]. *Ciona* releases gametes (oocytes and sperm) into sea water where fertilization and embryo development occur. These are processes very sensitive to substances present in the environment that come from industrial and agricultural activities and to chemical-physical variations of the environment (e.g., acidification, thermal elevation).

As filter feeders organisms, ascidians filter food from the incoming water and all the substances present in the environment are trapped on the pharyngeal mucus which is produced by the organ homologous to the mammalian thyroid. Consequently, ascidians are able to accumulate and concentrate harmful toxicants present in low concentrations in the surrounding water column [7]. Recently, it has been shown that ascidians are able to accumulate heavy metals such as vanadium and cadmium [8,9].

In this respect, a large body of literature reports the adverse impact of different type of stressors at the biochemical and cellular levels of ascidians inducing endocrine disruption and reproductive disorders with consequent fertilization success and reproductive fitness reduction and affecting in a long term even the survival of this cosmopolitan species [10-15].

Biomonitoring programs are aimed to evaluate the quality of the environment and require to identify model organisms able to function as bioindicators; these are defined as biological processes, species, or communities used to assess environmental toxicity [16].

The ideal bioindicator must be sessile for having not possibility to avoid the pollution source and filter-feeder since may process large amounts of water. In either cases they may provide a picture of environmental modifications as well as to identify polluted areas.

In the past, sea urchin has been considered the best bioindicator to evaluate environmental quality and, in particular, test on sperm and embryo development are from long time bioassays associated with monitoring [17,18]. Potential bioindicators to investigate pollutants impacts have been also recently proposed as molluscs [19], zebrafish [20] or others [21]. Since human is at the end of the food chain and human health is an end point for the research aimed to face up the threats of pollution and global change, there is evidence that deducing a human response to pollutants must come from comparing responses of organisms with similar physiology.

In our recent studies [22-24] we have provided evidences that *Ciona intestinalis* reproductive mechanisms are affected by either heavy metals or antifouling compounds. In particular, these xenobiotics exert an adverse impact on the gamete physiology, fertilization and larval development and morphology. The sensitivity to pollutants may partially explain the progressive reduction of *Ciona* populations observed in the last decades along the Neapolitan coast (personal communication).

Taken together these results are in favor to consider *Ciona intestinalis* a suitable bioindicator model due to its specific response to pollutants and its reproductive mechanisms may represent new bioassays for ecological risk assessment in the field of marine ecotoxicology (Figure 1).



Figure 1: *Ciona intestinalis*: from endangered species to top model.

*Corresponding author: Tosti E, Department of Biology and Evolution of Marine Organisms, Stazione Zoologica Anton Dohrn, Napoli, Italy, Tel: +390815833288; Fax: +390817641355; E-mail: tosti@szn.it

Received October 26, 2015; Accepted October 29, 2015; Published November 04, 2015

Citation: Gallo A, Tosti E (2015) The Ascidian *Ciona Intestinalis* as Model Organism for Ecotoxicological Bioassays. J Marine Sci Res Dev 5: e138. doi:10.4172/2155-9910.1000e138

Copyright: © 2015 Gallo A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Conclusion

Ascidians play a major role in the environment being cosmopolitan components of the marine ecosystems and species to protect. They have a peculiar position in evolution [25], in fact, due to the presence of a notochord at the larval stage and other features, are considered ancestors of chordates [26,27]. In consideration of the high representative value of *Ciona intestinalis* [28] among the marine invertebrates, the availability of its complete genome and the close phylogenetic relationship with vertebrates, the results herein reported substantiate that *Ciona* may be considered a new reliable model organism for ecotoxicology bioassays.

Acknowledgements

We thank Dr. Fiammetta Formisano for art work.

References

1. Satoh N (1994) Developmental biology of ascidians. Cambridge University Press, Cambridge.
2. Ciona Genome Group (2003) The *Ciona* genome issue. Dev Genes Evol 213: 1-218.
3. Tosti E, Dale B (1994) Regulation of fertilization current in ascidian oocytes by intracellular second messenger. Mol Reprod Dev 37: 473-476.
4. Tosti E, Gallo A, Silvestre F (2011) Ion currents involved in oocyte maturation, fertilization and early developmental stages of the ascidian *Ciona intestinalis*. Mol Reprod Dev 78: 854-860.
5. Russo GL, Kyojuka K, Antonazzo L, Tosti E, Dale B (1996) Maturation promoting factor in ascidian oocytes is regulated by different intracellular signals at meiosis I and II. Development 122: 1995-2003.
6. Russo GL, Bilotto S, Ciarcia G, Tosti E (2009) Phylogenetic conservation of Cytostatic Factor related genes in the Ascidian *Ciona intestinalis*. Gene 429: 104-111.
7. Villa L, Agati P, Mansueto C, Pellerito C, Scopelliti M, et al. (2003) Effects of tributyltin (IV) chloride on the gametes and fertilization of *Ascidia malaca* (Ascidacea: Tunicata). Appl Organomet Chem 17: 106-112.
8. Jiang A, Yu Z, Wang CH (2009) Bioaccumulation of cadmium in the ascidian *Styela clava* (Herdman 1881). African J Mar Sci 31: 289-295.
9. Treberg JR, Stacey JE, Driedzic WR (2012) Vanadium accumulation in ascidian coelomic cells is associated with enhanced pentose phosphate pathway capacity but not overall aerobic or anaerobic metabolism. Comp Biochem Physiol Part B Biochem Mol Biol 161: 323-330.
10. Bellas J (2006) Comparative toxicity of alternative antifouling biocides on embryos and larvae of marine invertebrates. Sci Tot Environ 367: 573-585.
11. Bellas J, Vázquez E, Beiras R (2001) Toxicity of Hg, Cu, Cd, and Cr on early developmental stages of *Ciona intestinalis* (Chordata, Ascidiacea) with potential application in marine water quality assessment. Water Res 35: 2905-2912.
12. Bellas J (2005) Toxicity assessment of the antifouling compound zinc pyrithione using early developmental stages of the ascidian *Ciona intestinalis*. Biofouling 21: 289-296.
13. Franchet C, Goudeau M, Goudeau H (1998) Tributyltin impedes early sperm-egg interactions at the egg coat level in the ascidian *Phallusia mammillata* but does not prevent sperm-egg fusion in naked eggs. Aquat Toxicol 44: 213-228.
14. Pennati R, Gropelli S, Zega G, Biggiogero M, De Bernardi F, et al. (2006) Toxic effects of two pesticides, Imazalil and Triadimefon, on the early development of the ascidian *Phallusia mammillata* (Chordata, Ascidiacea). Aquat Toxicol 79: 205-212.
15. Zega G, De Bernardi F, Gropelli S, Pennati R (2009) Effects of the azole fungicide Imazalil on the development of the ascidian *Ciona intestinalis* (Chordata, Tunicata): morphological and molecular characterization of the induced phenotype. Aquat Toxicol 91: 255-261.
16. Holt EA, Miller SW (2011) Bioindicators: using organisms to measure environmental impacts. Nature Educ.n Knowl. 3: 8.
17. Manzo S (2004) Sea urchin embryotoxicity test: proposal for a simplified bioassay. Ecotoxicol Environ Safety 57: 123-128.
18. Matranga V, Corsi I (2012) Toxic effects of engineered nanoparticles in the marine environment: model organisms and molecular approaches. Mar Environ Res 76: 32-40.
19. Waykar B, Deshmukh G (2012) Evaluation of bivalves as bioindicators of metal pollution in freshwater. Bull Environ Contam Toxicol 88: 48-53.
20. Segner H (2009) Zebrafish (*Danio rerio*) as a model organism for investigating endocrine disruption. Comp Biochem Physiol C 149: 187-195.
21. Zhou Q, Zhang J, Fu J, Shi J, Jiang G (2008) Biomonitoring: an appealing tool for assessment of metal pollution in the aquatic ecosystem. Anal Chim Acta 606: 135-150.
22. Gallo A, Silvestre F, Cuomo A, Papoff F, Tosti E (2011) The impact of metals on the reproductive mechanisms of the ascidian *Ciona intestinalis*. Mar Ecol 32: 222-231.
23. Gallo A, Tosti E (2013) Adverse effect of antifouling compounds on the reproductive mechanisms of the ascidian *Ciona intestinalis*. Mar Drugs 11: 3554-3568.
24. Gallo A, Tosti E (2015) Reprotoxicity of the antifoulant chlorothalonil in ascidians: an ecological risk assessment. Plos One 10: e0123074.
25. Kourakis MJ, Smith WC (2007) A conserved role for FGF signaling in chordate otic/atrial placode formation. Dev Biol 312: 245-257.
26. Delsuc F, Brinkmann H, Chourrout D, Philipp H (2006) Tunicates and not cephalochordates are the closest living relatives of vertebrates. Nature 439: 965-968.
27. Singh TR, Tsagkogeorga G, Delsuc F, Blanquart S, Shenkar N, et al. (2009) Tunicate mitogenomics and phylogenetics: peculiarities of the *Herdmania momus* mitochondrial genome and support for the new chordate phylogeny. BMC Genomics. 10: 534.
28. Sasakura Y, Inaba K, Satoh N, Kondo M, Akasaka K (2009) *Ciona intestinalis* and *Oxycomanthus japonicus*, representatives of marine invertebrates. Exp Anim 58: 459-469.