



The Essential Role of Physical Therapy in Post-Surgery Recovery

Giulia Rossi*

Department of Physiotherapy and Rehabilitation, Utrecht University, Netherlands

Introduction

Undergoing surgery, whether elective or necessary, is often a pivotal moment in a person's health journey. It represents a significant shift, as the body undergoes a physical change that requires time and care to heal. The immediate aftermath of surgery typically involves managing pain, monitoring for complications, and allowing the body to rest and recover. While these aspects are essential for initial recovery, one crucial element that significantly influences long-term outcomes is frequently overlooked: physical therapy.

Physical therapy is not merely a treatment to help patients "bounce back" from surgery; it is an integral part of the recovery process that can make the difference between a successful rehabilitation and a prolonged, painful recovery. Following surgery, the body may experience weakened muscles, reduced range of motion, and overall loss of function due to prolonged immobility or the physical trauma of the procedure itself. In these cases, physical therapy becomes vital in addressing these issues by providing targeted exercises and treatments designed to restore strength, improve flexibility, and enhance mobility [1].

Beyond just the physical aspects of recovery, physical therapy also helps reduce the risk of complications, such as blood clots, joint stiffness, or muscle atrophy. It also plays a key role in minimizing pain, potentially reducing the need for pain medications, including opioids, which carry their own set of risks and side effects. More importantly, physical therapy can significantly speed up recovery, empowering patients to regain their independence, return to their daily activities, and improve their overall quality of life.

In this article, we will explore how physical therapy contributes to post-surgery recovery by detailing its benefits, the processes involved, and why it should be a fundamental part of every post-operative care plan. Whether recovering from a joint replacement, spinal surgery, or soft tissue repair, physical therapy is an essential tool that helps patients rebuild their bodies and minds after surgery, ultimately guiding them towards a healthier, more active life [2].

The role of physical therapy in post-surgery recovery

Physical therapy in the post-surgery phase focuses on rehabilitation to restore movement and function to the body. Its role extends far beyond just helping a patient "get moving" again. PT is tailored to address the unique needs of each individual depending on the type of surgery, the patient's overall health, and their long-term goals. The primary goals of physical therapy in this context are to reduce pain, promote healing, prevent complications, and improve physical function [3].

Pain management

Pain is an inevitable part of the post-surgical experience, and managing it is essential to a successful recovery. Physical therapists utilize various techniques such as manual therapy, soft tissue mobilization, therapeutic exercises, and modalities like heat, cold, or electrical stimulation to alleviate discomfort [4]. By focusing on the

areas affected by the surgery, physical therapy can help reduce pain and limit reliance on medications, particularly opioids, which can carry risks for addiction or side effects.

Restoring mobility

After surgery, patients often experience stiffness, weakness, and limited range of motion. For example, after a knee or hip replacement, many people struggle to move their joints properly due to scar tissue, muscle atrophy, or pain. Physical therapists design specific exercises that target the injured or operated area to rebuild strength, improve flexibility, and restore joint function. By working on mobility, physical therapy helps patients regain their independence in activities of daily living, such as walking, dressing, or climbing stairs.

Preventing complications

A critical aspect of post-surgical rehabilitation is preventing complications such as blood clots, joint stiffness, or muscle atrophy. For instance, patients recovering from orthopedic surgeries like joint replacements or spinal fusions are at risk of developing blood clots if they remain immobile for too long. Physical therapists guide patients through gentle movements and exercises that stimulate circulation and reduce the risk of such complications. In addition, they can teach patients proper body mechanics to avoid strain on vulnerable parts of the body during recovery [5].

Building strength and endurance

Following surgery, it's common for muscles to weaken or shrink, particularly if the patient has been immobile for an extended period. Physical therapy focuses on rebuilding muscle strength and endurance through progressive resistance exercises. Strengthening the muscles surrounding the surgical site not only aids in faster recovery but also protects the joint or area from further injury [6]. In many cases, physical therapy involves exercises that improve balance, stability, and coordination, helping the patient return to their pre-surgery level of function.

Psychological support and motivation

A comprehensive post-surgical recovery plan goes beyond the physical aspects and addresses the emotional challenges patients face. Surgery, particularly major procedures, can leave patients feeling

*Corresponding author: Giulia Rossi, Department of Physiotherapy and Rehabilitation, Utrecht University, Netherlands, E-mail: Giulia.r@hotmail.com

Received: 02-Nov-2024, Manuscript No: jnp-24-154387; Editor assigned: 04-Nov-2024, Pre-QC No: jnp-24-154387(PQ); Reviewed: 18-Nov-2024, QC No: jnp-24-154387; Revised: 23-Nov-2024, Manuscript No: jnp-24-154387(R); Published: 30-Nov-2024, DOI: 10.4172/2165-7025.1000769

Citation: Giulia R (2024) The Essential Role of Physical Therapy in Post-Surgery Recovery. J Nov Physiother 14: 769.

Copyright: © 2024 Giulia R. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

frustrated, anxious, or demoralized as they work through recovery [7]. Physical therapists offer encouragement, set achievable goals, and track progress. This holistic approach boosts patients' confidence and motivates them to stay committed to their rehabilitation program [8].

Conclusion

Physical therapy is a vital component of post-surgery recovery, offering benefits that go far beyond simple rehabilitation. It not only accelerates healing by addressing pain, restoring mobility, and preventing complications, but it also plays an integral role in rebuilding strength and improving overall functionality. For patients recovering from surgery, working with a skilled physical therapist can make the difference between a slow, painful recovery and a faster, more complete return to normal life. By incorporating physical therapy into the post-operative care plan, individuals can significantly improve their chances of a full recovery and regain their independence more swiftly.

Whether recovering from joint surgery, spinal surgery, or any other procedure, patients should recognize the value of physical therapy as an essential partner in their recovery process. By following a personalized rehabilitation plan, patients can maximize their recovery potential and enjoy a better quality of life in the long term.

Acknowledgement

None

Conflict of Interest

None

References

1. Sackett D, Rosenberg WM, Gray JA, Haynes RB, Richardson WS (1996) Evidence based medicine: What it is and what it isn't. *Br Med J* 312: 71-72.
2. Payton O, Nelson CE (1996) A preliminary study of patients perceptions of certain aspects of their physical therapy experience. *Physiother Theory Pract* 12: 27-38.
3. Lerner-Frankiel M, Vargas S, Brown M, Krusell L, Schoneberger W (1986) Functional community ambulation: What are your criteria?. *Clin Manage Phys Ther* 6: 12-15.
4. Stephens JM, Goldie PA (1999) Walking speed on parquetry and carpet after stroke: effect of surface and retest reliability. *Clin Rehabil* 13: 171-181.
5. Wade DT, Wood VA, Heller A, Maggs J (1987) Walking after stroke: Measurement and recovery over the first 3 months. *Scand J Rehabil Med* 19: 25-30.
6. Perry J, Garrett M, Gronley J, Mulroy S (1995) Classification of walking handicap in the stroke population. *Stroke* 26: 982-989.
7. Bernhardt J, Ellis P, Denisenko S, Hill K (1998) Changes in balance and locomotion measures during rehabilitation following stroke. *Physiother Res Int* 3: 109-122.
8. H0ill K, Schwarz J, Kalogeropoulos A, Gibson S (1996) Fear of falling revisited. *Arch Phys Med Rehabil* 77: 1025-1029.