



# The Science Behind Weight Loss: Understanding Metabolism in Morbid Obesity

Priyanka Sharma\*

Department of Biotechnology, Kalinga Institute of Industrial Technology, India

## Introduction

Weight loss is often oversimplified as a matter of “calories in versus calories out.” While this basic principle holds some truth, the science behind weight loss especially in the context of morbid obesity is far more complex. Metabolism, the process by which the body converts food into energy, plays a critical role in weight management. For individuals with morbid obesity (defined as a Body Mass Index, or BMI, of 40 or higher), metabolic factors contribute significantly to the challenges of losing weight. Metabolic rates, hormonal imbalances, and adaptive responses to weight loss create a unique and often frustrating barrier to achieving sustainable results [1]. This article delves into the science behind metabolism in morbid obesity, offering insight into why losing weight is more difficult for some and what strategies can help overcome these challenges.

## Description

### Understanding metabolism and its role in weight management

**Basal metabolic rate (BMR):** Metabolism can be broken down into several components, with Basal Metabolic Rate (BMR) being the most significant. BMR refers to the number of calories the body needs to maintain basic physiological functions such as breathing, circulation, and cell production while at rest. It accounts for 60-70% of the total energy expenditure in most people. For individuals with morbid obesity, BMR is often higher because the body needs more energy to support the extra weight. However, this does not necessarily make weight loss easier, as metabolic adaptations often counteract efforts to reduce body mass [2].

**Adaptive thermogenesis:** When people with morbid obesity attempt to lose weight, their bodies undergo a process known as adaptive thermogenesis. This is the body’s way of conserving energy in response to reduced caloric intake. Essentially, the body becomes more efficient, burning fewer calories to perform the same tasks. This phenomenon makes it difficult to continue losing weight after an initial period of success, commonly referred to as a “weight loss plateau.” The body perceives weight loss as a threat and tries to preserve energy by slowing down metabolism, a survival mechanism that evolved during times of food scarcity [3].

**Hormonal influences on metabolism:** Hormones play a crucial role in regulating appetite, fat storage, and energy expenditure. In individuals with morbid obesity, hormonal imbalances can significantly affect metabolism. For instance, leptin is a hormone that signals the brain to stop eating when fat stores are sufficient. However, in people with morbid obesity, leptin resistance can occur, meaning the brain no longer responds appropriately to leptin signals, leading to overeating and reduced energy expenditure [4]. Similarly, insulin, the hormone responsible for regulating blood sugar, may become less effective in people with morbid obesity, contributing to fat storage and weight gain.

Additionally, levels of the hormone ghrelin, which stimulates hunger, often rise during weight loss efforts, making it more

challenging to maintain caloric restriction. These hormonal changes make it difficult for individuals with morbid obesity to sustain long-term weight loss, as their bodies fight to regain lost weight.

### The challenges of weight loss in morbid obesity

**Metabolic slowdown:** As weight decreases, so does the body’s need for energy. This natural reduction in calorie requirements, combined with adaptive thermogenesis, can lead to a significant metabolic slowdown. For example, someone who loses a substantial amount of weight may find that they burn far fewer calories than someone of the same weight who has never been obese [5]. This creates an ongoing challenge, as maintaining weight loss requires consuming fewer calories than the body burns, often to an extreme degree.

**Loss of lean muscle mass:** Another key factor in the difficulty of weight loss in morbid obesity is the loss of lean muscle mass. Muscle is metabolically active, meaning it burns more calories than fat, even at rest. During weight loss, particularly rapid or extreme weight loss, both fat and muscle mass are lost. The reduction in muscle mass further slows metabolism, making it even harder to maintain weight loss over time. Strength training and resistance exercises are essential to minimize muscle loss and help preserve metabolic function during weight loss efforts.

**Weight regain and the set point theory:** Many individuals with morbid obesity experience weight regain after initial success. This can be attributed to a combination of metabolic slowdown, hormonal changes, and the body’s attempt to return to a “set point.” The set point theory suggests that the body has a natural weight range that it defends, making it difficult to deviate from this range for extended periods. For those with morbid obesity, the set point may be higher than what is considered a healthy weight, making it particularly challenging to lose weight and maintain the loss [6].

### Strategies to support weight loss in morbid obesity

**Gradual weight loss:** One key strategy for overcoming metabolic challenges is gradual weight loss. Rapid weight loss often triggers more pronounced metabolic adaptations, whereas slower, steady weight loss allows the body to adjust with fewer defensive responses. A combination of moderate caloric restriction, regular physical activity, and behavioral changes can help achieve sustainable results.

\*Corresponding author: Priyanka Sharma, Department of Biotechnology, Kalinga Institute of Industrial Technology, India, E-mail: priya\_sh@gmail.com

**Received:** 03-Sep-2024, Manuscript No: jowt-24-149876, **Editor assigned:** 05-Sep-2024, Pre QC No: jowt-24-149876(PQ), **Reviewed:** 19-Sep-2024, QC No: jowt-24-149876, **Revised:** 23-Sep-2024, Manuscript No: jowt-24-149876(R) **Published:** 30-Sep-2024, DOI: 10.4172/2165-7904.1000730

**Citation:** Priyanka S (2024) The Science Behind Weight Loss: Understanding Metabolism in Morbid Obesity. J Obes Weight Loss Ther 14: 730.

**Copyright:** © 2024 Priyanka S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Incorporating resistance training:** As mentioned earlier, preserving muscle mass is crucial for maintaining metabolism during weight loss. Incorporating resistance training or strength exercises into a weight loss program can help minimize muscle loss and support long-term weight management. This approach not only aids in fat loss but also helps maintain lean body mass, keeping the metabolism more active [7].

**Addressing hormonal imbalances:** In some cases, medical intervention may be necessary to address underlying hormonal issues that complicate weight loss. For example, managing insulin resistance through dietary changes or medication can help reduce fat storage and support more effective weight management. Leptin resistance, though harder to target directly, can be improved through gradual weight loss and a focus on overall health rather than just rapid results.

**Metabolic surgery:** For individuals with morbid obesity who have struggled to lose weight through traditional means, metabolic surgery, such as gastric bypass or sleeve gastrectomy, may offer a more effective option. These surgeries not only reduce the size of the stomach but also have profound effects on gut hormones that regulate hunger and metabolism [8]. By altering the digestive system, metabolic surgery can help reset the body's weight-regulating mechanisms, making it easier to lose and maintain weight.

## Conclusion

Understanding the science behind metabolism is essential for appreciating the unique challenges of weight loss in individuals with morbid obesity. Metabolic adaptations, hormonal imbalances, and the body's natural tendency to resist weight loss make this journey more difficult than a simple equation of calories in versus calories out. However, with a combination of gradual weight loss strategies,

resistance training, and, in some cases, medical intervention, it is possible to overcome these barriers. By addressing both the metabolic and behavioral aspects of morbid obesity, individuals can achieve sustainable weight loss and improve their overall health.

## Acknowledgement

None

## Conflict of Interest

None

## References

1. Fonesca H, Matos MG, Guerra A, Pedro JG (2009) Are overweight and obese adolescents different from their peers. *Int J Pediatr Obes* 4: 166-174.
2. Freedman DS, Khan LK, Dietz WH, Srinivasan SR, Berenson GS (2001) Relationship of childhood obesity to coronary heart disease risk factors in adulthood The Bogalusa Heart Study. *Pediatrics* 108: 712-718.
3. Huelsing J, Kanafani N, Mao J, White NH (2010) Camp Jump Start: effects of a residential summer weight-loss camp for older children and adolescents. *Pediatrics* 125: 884-890.
4. l'Allemand-Jander D (2010) Clinical diagnosis of metabolic and cardiovascular risks in overweight children: early development of chronic diseases in the obese child. *Int J Obes* 34 Suppl 2: S32-36.
5. Castagneto M, De Gaetano A, Mingrone G, Tacchino R, Nanni G, et al. (1994) Normalization of insulin sensitivity in the obese patient after stable weight reduction with biliopancreatic diversion. *Obes Surg* 4: 161-168.
6. Cooper C, Sarvey S, Collier D, Willson C, Green I, et al. (2006) For comparison: experience with a children's obesity camp. *Surg Obes Relat Dis* 2: 622-626.
7. Cowan GS Jr, Buffington CK (1998) Significant changes in blood pressure, glucose, and lipids with gastric bypass surgery. *World J Surg* 22: 987-992.
8. Fagot-Champagna A, Pettit DJ, Engelgau MM, Burrows NR, Geiss LS, et al. (2000) Type 2 Diabetes among North American children and adolescents: An epidemiologic review and a public health perspective. *J Pediatr* 136: 664-672.