

## Use of LAB to Control Urogenital Infections

Mittu B\*, Kaur B and Balgir PP

Department of Biotechnology, Punjabi University, Patiala, India

\*Corresponding author: Mittu B, Department of Biotechnology, Punjabi University, Patiala, India, E-mail: bharti9mittu@yahoo.com

Received date: November 3, 2015; Accepted date: January 8, 2016; Published date: January 16, 2016

Copyright: © 2016 Mittu, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

### Introduction

A urinary tract infection (UTI) is an **infection** that affects part of the **urinary tract**. When it affects the lower urinary tract, it is known as a simple **cystitis** (a bladder infection) and when it affects the upper urinary tract, it is known as **pyelonephritis** (a kidney infection). The indigenous microbiota plays an important role in protecting the host from colonization of invading pathogens. *Lactobacillus* is the predominant genus in the vaginal [1] and endocervical microbial communities [2] and is present at concentrations of 10<sup>7</sup> to 10<sup>8</sup> CFU/ml of vaginal fluid in healthy postmenopausal/ premenopausal women [3].

There is an emerging role of lactobacilli in the control and maintenance of the vaginal bacterial microflora. *L. jensenii* and *L. gasseri* are two of the most common species present, as determined by culture-independent techniques [4]. Relationship between LAB and vaginal pathogen have been studied which indicates that antibiotic treatment should not kill healthy LAB to preserve healthy vagina [4]. Pascual et al. (2010) found that *L. fermentum* L23 isolated from vaginal swabs of healthy woman can prevent and cure *E. coli* infection in a murine vaginal tract model [5].

*Lactobacillus* sp. ferment glycogen secreted by vaginal epithelial cells into lactic acid, and colonization by these microorganisms

correlates to the low pH in the vagina [6,7]. The acidic environment protects the vagina against over-growth of pathogens [8,9]. *L. salivarius* subsp. *salivarius* CRL 1328 was able to inhibit the growth of *E. coli*, *Klebsiella* sp., *G. vaginalis*, *S. aureus* and *S. agalactiae* by the effect of pH, and *N. gonorrhoeae* and *E. faecalis* by a bacteriocin-like substance that was reported previously [10-12]. *L. acidophilus*, *L. brevis*, *L. plantarum*, *L. casei*, *L. lactis* subsp. *lactis*, *L. raffinolactis*, *P. acidilactici* and *P. pentosaceus* isolated from healthy female vaginal fluid, are able to inhibit the growth of *C. albicans*, *G. vaginalis* and *N. gonorrhoeae*, *Escherichia coli*, *C. albicans* and *S. aureus* [13-17].

Earlier, Raiz and others (2010) isolated 72 strains from yogurt and fecal materials of human, chick, parrot and cat. Only two isolates namely *L. fermentum* and *L. acidophilus* were found to produce bacteriocins having antimicrobial potential against cephalosporin resistant *E. coli*. In a clinical trial performed using capsules containing *L. fermentum* RC-14 and *L. rhamnosus* GR-1, Gil et al. (2010) have proven their effectiveness for the treatment of patients with vulvovaginal candidiasis. Recently, *L. fermentum* was isolated from human milk and its bacteriocin has been assessed for preventing urinary tract infections [18-23]. Probiotics recommended for eradication of common human UTI vaginal pathogens are enlisted in Table A [24-26].

Causal Organism	Disease	Symptoms	Recommended Probiotics	References
<i>Candida albicans</i>	Vaginitis, Vulvovaginal candidiasis	Irritation and soreness of the vulva, a thick, white vaginal discharge that doesn't usually smell.	<i>L. pentosus</i> TV35b	[19]
<i>Enterococcus faecalis</i>	UTI	Blood in urine, pyelonephritis	<i>L. salivarius</i> subsp. <i>salivarius</i> CRL 1328	[10]
<i>Enterococcus faecium</i>	UTI	Normal commensals in human vagina, but if exceeds in counts can cause disease.	<i>L. salivarius</i> subsp. <i>salivarius</i> CRL 1328	[11]
<i>Escherichia Coli</i>	UTI cystitis	Pain or burning with urination, lower abdominal pain or pressure, and/or the need to urinate frequently; cloudy darker or bloody urine	<i>L. acidophilus</i> CRL 1259, <i>L. crispatus</i> 21L07	[24,25]
			<i>L. jensenii</i> 5L08	[17]
<i>Escherichia Coli</i>	UTI cystitis	Pain or burning with urination, lower abdominal pain or pressure, and/or the need to urinate frequently; cloudy darker or bloody urine	<i>L. acidophilus</i> CRL 1259, <i>L. crispatus</i> 21L07	[24,25]
			<i>L. jensenii</i> 5L08	[17]

<i>Klebsiella pneumonia</i>	UTI	Burning sensation during urination, back pain, abdominal pain	<i>L. salivarius</i> subsp. <i>salivarius</i> CRL 1328	[21]
<i>Neisseria gonorrhoeae</i>	Gonorrhoea	Vaginal discharge and pain during urination, urethritis, vulvovaginitis	<i>L. salivarius</i> subsp. <i>salivarius</i> CRL 1328	[10,11]
			<i>L. jenseni</i> ATCC 25258, <i>L. gasseri</i> ATCC 33323	[21]
<i>Pseudomonas aeruginosa</i>	Burning sensation during urination, back pain, abdominal pain	Hospital-acquired urinary tract infections from catheterization, instrumentation or surgery	<i>L. gasseri</i> TL093C, TL143A, TL029, TL080 <i>L. acidophilus</i> TL099 <i>L. delbrueckii</i> TL059	[15]
<i>Staphylococcus aureus</i>	Toxic shock syndrome	Variety of local and systemic infections	<i>L. bulgaricus</i>	[14]
			<i>L. paracasei</i> CRL 1289	[26]
<i>Streptococcus agalactiae</i>	UTI	Bacteria can be passed from a pregnant woman to her baby during labor, if she is a carrier of the bacteria.	<i>L. bulgaricus</i>	[24,25]

**Table A:** Common Opportunistic Pathogens of Human Urogenital System and their control by probiotic LAB.

**References**

1. Andreu A (2004) Lactobacillus as a probiotic for preventing urogenital infections. Rev Med Microbiol 15: 1-6.
2. Wilson RD, Fricke PM, Leibfried-Rutledge ML, Rutledge JJ, Penfield CM, et al. (2006) In vitro production of bovine embryos using sex-sorted sperm. Theriogenology 65: 1007-1015.
3. Redondo-Lopez V, Cook RL, Sobel JD (1990) Emerging role of lactobacilli in the control and maintenance of the vaginal bacterial microflora. Rev Infect Dis 12: 856-872.
4. Razzak MS, Al-Charrakh AH, Al-Greitty BH (2011) Relationship between lactobacilli and opportunistic bacterial pathogens associated with vaginitis. N Am J Med Sci 3: 185-192.
5. Pavlova SI, Kilic AO, Kilic SS, So JS, Nader-Macias ME, et al. (2002) Genetic diversity of vaginal lactobacilli from women in different countries based on 16S rRNA gene sequences. J Appl Microbiol 92: 451-459.
6. Boskey ER, Cone RA, Whaley KJ, Moench TR (2001) Origins of vaginal acidity: high D/L lactate ratio is consistent with bacteria being the primary source. Hum Reprod 16: 1809-1813.
7. Rönnqvist PD, Forsgren-Brusk UB, Grahn-Håkansson EE (2006) Lactobacilli in the female genital tract in relation to other genital microbes and vaginal pH. Acta Obstet Gynecol Scand 85: 726-735.
8. Martin LS, McDougal JS, Loskoski SL (1985) Disinfection and inactivation of the human T lymphotropic virus type III/Lymphadenopathy-associated virus. J Infect Dis 152: 400-403.
9. Hill JA, Anderson DJ (1992) Human vaginal leukocytes and the effects of vaginal fluid on lymphocyte and macrophage defense functions. Am J Obstet Gynecol 166: 720-726.
10. Ocaña VS, de Ruiz Holgado AA, Nader-Macias ME (1999) Growth inhibition of Staphylococcus aureus by H2O2-producing Lactobacillus paracasei subsp. paracasei isolated from the human vagina. FEMS Immunol Med Microbiol 23: 87-92.
11. Ocaña VS, Pesce De Ruiz Holgado AA, Nader-Macias ME (1999) Characterization of a bacteriocin-like substance produced by a vaginal Lactobacillus salivarius strain. Appl Environ Microbiol 65: 5631-5635.
12. Ocaña VS, Pesce De Ruiz Holgado AA, Nader-Macias ME (1999) Characterization of a bacteriocin-like substance produced by a vaginal Lactobacillus salivarius strain. Appl Environ Microbiol 65: 5631-5635.
13. Selvamohan T, Sujitha S (2010) Antimicrobial activity of a probiotic Lactobacillus Plantarum against urinary tract infection (UTI) causing pathogens. J Der Pharmacia Lettre 2: 432-440.
14. Amin M, Goodarzi H, Orang Z, Farsi S, Jorfi M (2011) Isolation and identification of Lactobacillus species from the vagina and their antimicrobial properties. Afr. J. Microbiol. Res 5: 3300-3304.
15. Alpay-Karaoglu, Aydin SE, Kilic SS, Kilic AO (2003) Antimicrobial activity and characteristics of bacteriocins produced by vaginal lactobacilli. Turk J Med Sci 33: 7-12.
16. Gil NE, Martinez RC, Gomes BC, Nomizo A, De Martinis EC (2010) Vaginal lactobacilli as potential probiotics against Candida SPP. Braz J Microbiol 41: 6-14.
17. Kaewsrichan J, Peeyananjarasri K, Kongprasertkit J (2006) Selection and identification of anaerobic lactobacilli producing inhibitory compounds against vaginal pathogens. FEMS Immunol Med Microbiol 48: 75-83.
18. Ilayaraja R, Radhamadhavan G (2011) Assessment for potential use of bacteriocin producing Lactobacillus fermentum isolated from human milk for preventing urinary tract infections. J.Pharmacy Res 4: 3445-3447.
19. Okkers DJ, Dicks LM, Silvester M, Joubert JJ, Odendaal HJ (1999) Characterization of pentocin TV35b, a bacteriocin-like peptide isolated from Lactobacillus pentosus with a fungistatic effect on Candida albicans. J Appl Microbiol 87: 726-734.
20. Pascual L, Ruiz F, Giordano W, Barberis IL (2010) Vaginal colonization and activity of the probiotic bacterium Lactobacillus fermentum L23 in a murine model of vaginal tract infection. J Med Microbiol 59: 360-364.
21. St Amant DC, Valentin-Bon IE, Jerse AE (2002) Inhibition of Neisseria gonorrhoeae by Lactobacillus species that are commonly isolated from the female genital tract. Infect Immun 70: 7169-7171.
22. Saunders S, Bocking A, Challis J, Reid G (2007) Effect of Lactobacillus challenge on Gardnerella vaginalis biofilms. Colloids Surf B Biointerfaces 55: 138-142.
23. Riaz S, Kashif Nawaz S, Hasnain S (2010) Bacteriocins produced by L. Fermentum and L. Acidophilus can inhibit cephalosporin resistant e. Coli. Braz J Microbiol 41: 643-648.

- 
24. Juárez Tomás MS, Ocaña VS, Wiese B, Nader-Macías ME (2003) Growth and lactic acid production by vaginal *Lactobacillus acidophilus* CRL 1259, and inhibition of uropathogenic *Escherichia coli*. *J Med Microbiol* 52: 1117-1124.
  25. Juárez Tomás MS, Ocaña VS, Wiese B, Nader-Macías ME (2003) Growth and lactic acid production by vaginal *Lactobacillus acidophilus* CRL 1259, and inhibition of uropathogenic *Escherichia coli*. *J Med Microbiol* 52: 1117-1124.
  26. Zarate G, Santos V, Nader E (2007) Protective effect of vaginal *Lactobacillus paracasei* CRL 1289 against urogenital infection produced by *Staphylococcus aureus* in a mouse animal model. *Inf Dis Obstet Gynecol*: 48358.