

## Role of Heat Shock Proteins in Alzheimer's Disease

Ubaldo Armato\*

Department of Life and Reproduction Sciences, University of Verona Medical School, Verona, Italy

\*Corresponding author: Ubaldo Armato, Department of Life and Reproduction Sciences, University of Verona Medical School, Verona, Italy, E-mail: ubaldo.armato@univr.it

Received: 5-Jan-2022, Manuscript No. JADP-21-55842; Editor assigned: 7-Jan-2022, PreQC No. JADP-21-55842 (PQ); Reviewed: 21-Jan-2022, QC No. JADP-21-55842; Revised: 25-Jan-2022, Manuscript No. JADP-21-55842 (R); Published: 1-Feb-2022, DOI: 10.4172/2161-0460.1000532.

Citation: Armato U (2022) Role of Heat Shock Proteins in Alzheimer's Disease. J Alzheimers Dis Parkinsonism 12: 532.

Copyright: © 2022 Armato U. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Description

Most age-related neurodegenerative diseases are hallmarked by the existence of pathological protein depositions in the brain, for example, extracellular plaques and neurofibrillary tangles in Alzheimer's disease. Furthermore, aging and neurodegeneration in Alzheimer's disease are often accompanied by a functionally impaired UPS that may be both a cause and one of the detrimental consequences of the aggregation. Synaptic plasticity is affected in Alzheimer's disease and involved metabolism of at least 5%-8% of brain proteins each day. As stated, central to PQC are HSPs that interact with mis or unfolded proteins. They generally bind to exposed hydrophobic, aggregation-prone regions, preventing their self-association and aggregation. Whilst the best-known outcome of such interaction is that chaperone next assists in the refolding of its bound client protein, HSPs also can assist in supporting its client's proteosomal or autophagosomal degradation. In addition, HSPs can assist in the disassembly of protein complexes and provide the driving force for translocation of proteins across membranes.

From gene transcription to posttranslational modification, several quality checks are essential before a protein is ready for biological action, locally or distantly *via* axonal transport in neurons. The accurate folding and control over levels of many proteins must be tightly regulated both spatially and temporally. To achieve this, the cell possesses a network of different protein quality control systems for cotranslational protein folding *via* molecular chaperons, as the first line of defense against protein misfolding and aggregation. Subsequently, accurate protein degradation, encompassing the ubiquitin-proteasome system and the autophagosomal-lysosomal system is the second line defense. Recent data also suggest an additional protein quality control pathway in which misfolded proteins are excreted actively after encapsulation at the endoplasmic reticulum, a process dubbed Misfolded Associated Protein Secretion.

In each of the protein quality control systems, an array of Heat Shock Proteins (HSPs) are required to chaperone and direct substrates toward their correct fate and to avoid off-pathway reactions that might lead to harmful accumulations of protein aggregates. Recent studies focused on the cross-talk between the UPS and the HSPs, concepts that are in line with the current idea that Alzheimer's disease is in an asymptomatic cellular phase long before it can be translated into clinically relevant conservation and interventions.

As stated before, recent data suggest an additional PQC pathway in which misfolded proteins are excreted actively after encapsulation at the endoplasmic reticulum, a process dubbed MAPS. This process was

dependent on an ER-resident ubiquitin-specific protease, USP19, which originally was shown to function in ER-associated proteosomal degradation. The catalytic domain of USP19 was found to possess an unprecedented chaperone activity, allowing recruitment of misfolded protein to the ER surface. Deubiquitinated cargos are then encapsulated into ER-associated late endosomes and secreted to the cell exterior. The action of USP19 was found to be independent of its interaction with HSP90, with whom it normally partners in the process of ER-associated proteosomal degradation. In a recent research, extracellular release of neurodegenerative-associated proteins was also found to be a regulated process, operating likely under conditions of compromised intracellular PQC. Finally a recent study on *C. elegans* showed that adult neurons extrude large membrane-surrounded vesicles, termed exospheres. This effect was likewise augmented upon decline in intercellular PQC capacity as well as by compressed mitochondrial functioning. Although each of these export mechanisms may provide a level of cytoprotection, this type of secretion may also contribute to the prion like propagation of A $\beta$  or tau and therefore may be a target for inhibition to halt disease progression. These studies also further underscore the importance of maintaining an optimal intracellular PQC to prevent neurodegenerative diseases and brain aging.

Protein homeostasis is crucial for the proper function of cells and maintaining an optimally functioning PQC network is therefore essential to prevent neuronal degeneration. Most facets of PQC erode during aging and this may be a central factor in the onset or progression of Alzheimer's disease. Conversely, potentiating PQC in various ways may help to delay the disease onset, either directly *via* various means of chaperoning A $\beta$  or tau monomers or aggregates or indirectly by maintaining normal PQC and preventing aggregate release and propagation. A summary of possible actions of HSP in the pathogenesis of Alzheimer's disease is provided as a result of the research in the recent years. Simultaneously, improving UPS efficiency *via* DUBs or autophagic activities is an option. Alternatively, reducing the burden on the PQC, e.g., by preventing accumulation of other misfolded proteins in general or even specific ones like UBB<sup>+1</sup> could also help to delay a disruption in protein homeostasis. However as many of these processes are promiscuous, interference with their global activity may come with too many negative tradeoffs. They may also only be effective transiently due to auto regulatory feedback loops or epigenetic changes, as observed when trying to boost HSF-1 actively by HSP90 inhibition. Therefore, specific targeting of cochaperons or UBB<sup>+1</sup> or their transcripts may be an option.