



Weight Loss Doesn't Help Pregnancy Chances, Study Finds

Joe Thomas*

Jawaharlal Nehru University, Hyderabad, India

Introduction

Women who are obese and struggling to become pregnant are often advised to lose weight, but a new study finds no fertility benefits from weight loss.

A randomized study of 379 women with obesity and unexplained infertility found that intensive lifestyle changes that shed pounds led to no better chances of pregnancy and healthy births than simply increasing physical activity without weight loss [1].

"We have known for decades that obese women often have difficulty getting pregnant, said researcher. For this reason, many physicians advise weight loss prior to conception. However, there are few studies that have addressed the issue comparing a healthy lifestyle i.e., exercise vs. exercise plus weight loss.

Description

Obesity and Pregnancy

The FIT-PLESE study, conducted at nine academic medical centers across the country, divided participants into two groups: Half the women dieted intensely using meal replacements, medications and increased physical activity. The other half simply increased their physical activity without trying to lose weight. After completing the programs, both groups received three rounds of standard infertility treatments [2].

Women in the weight-loss program ended up losing, on average, 7% of their body weight, while participants in the exercise-only group typically maintained their weights. But, in the end, there were no significant differences between the two groups in terms of the frequency of healthy births. In total, 23 of the 188 women who completed the 16-week intensive weight-loss program ended up giving birth; among the 191 who completed the exercise-only program, 29 gave birth [3].

The intensive dieting program did offer health benefits for the women who completed it, however. In addition to dropping pounds, they saw a major decrease in metabolic syndrome, a cluster of conditions that increase the risk for serious health problems such as diabetes, stroke and heart disease.

Based on their findings, researchers conclude that the weight-loss program did not make women more fertile or improve birth outcomes compared with simply exercising. They note the health benefits of weight loss may not translate into better odds of getting pregnant [4].

Weight loss improved metabolic health in these subjects. Unfortunately the changes seen did not improve fertility. Infertility within this population remains an important health issue, and will require further studies to address the problem in the future.

Health care professionals often encourage women with obesity to lose weight prior to trying to conceive or start infertility treatments [5]. But a new nationwide study led by Penn State College of Medicine found that women with obesity and unexplained infertility who lost weight prior to starting infertility treatments did not have a greater chance of having a healthy baby than those who did not lose weight prior to starting therapy.

Conclusion

40% of U.S. women between the ages of 20 and 40 are estimated to have obesity, which has been associated with trouble getting pregnant, pregnancy complications and loss. As a result, it is common for those women to receive guidance to lose weight prior to conception to increase their chances of having a healthy baby.

Acknowledgement

None

Conflict of Interest

None

References

1. Paul RW, Elder L, Bartell T (1997) California teacher preparation for instruction in critical thinking: Research findings and policy recommendations.
2. Farnum J (2005) Sense of place in natural resource recreation and tourism: An evaluation and assessment of research findings.
3. Thornton PL, Kieffer EC, Salabarría-Peña Y, Odoms-Young A, Willis SK, et al. (2006) Weight, diet, and physical activity-related beliefs and practices among pregnant and postpartum Latino women: the role of social support. *Matern Child Health J* 10(1): 95-104.
4. Rose DC, Sutherland WJ, Parker C, Lobley M, Winter M, et al. (2016) Decision support tools for agriculture: Towards effective design and delivery. *Agric Syst* 149: 165-174.
5. Sutton RI, Hargadon A (1996) Brainstorming groups in context: Effectiveness in a product design firm. *Administrative science quarterly* pp: 685-718.

*Corresponding author: Joe Thomas, Jawaharlal Nehru University, Hyderabad, India, E-mail: joe.thomas@gmail.com

Received: 03-May-2022, Manuscript No. JOWT-22-498; **Editor assigned:** 04-May-2022, PreQC No. JOWT-22-498(PQ); **Reviewed:** 18-May-2022, QC No. JOWT-22-498; **Revised:** 23-May-2022, Manuscript No. JOWT-22-498(R); **Published:** 30-May-2022, DOI: 10.4172/2165-7904.1000498

Citation: Thomas J (2022) Obesity More than Obstructive Sleep Apnea Plays a Crucial Role to Increase the Cardiovascular Risk. *J Obes Weight Loss Ther* 12: 498.

Copyright: © 2022 Thomas J. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.